

## Methods of Treating Obesity Using A Neurotensin Receptor Ligand

Cross-Reference to Related Invention

This application claims priority of U.S. provisional application number  
5 60/199,951, filed April 27, 2000.

Field of the Invention

The present invention relates to methods of treating obesity, diabetes, sexual  
dysfunction (including erectile dysfunction), atherosclerosis, insulin resistance,  
impaired glucose tolerance, hypercholesterolemia, or hypertriglyceridemia using a  
10 compound that is a neurotensin receptor ligand. The present invention also relates to  
compositions and kits that comprise a neurotensin receptor ligand.

Background of the Invention

Obesity is a devastating disease. In addition to harming physical health,  
15 obesity can wreak havoc on mental health because obesity affects self-esteem,  
which ultimately can affect a person's ability to interact socially with others.  
Unfortunately, obesity is not well understood, and societal stereotypes and  
presumptions regarding obesity only tend to exacerbate the psychological effects of  
the disease. Because of the impact of obesity on individuals and society, much effort  
20 has been expended to find ways to treat obesity, but little success has been achieved  
in the long-term treatment and/or prevention of obesity.

Neurotensin is a thirteen amino acid peptide that appears to have functions as  
a neurotransmitter and neuromodulator in the nervous system and as a local  
hormone in the periphery. Specifically, neurotensin is a neuromodulator of dopamine  
25 transmission and of anterior pituitary hormone secretion, and exerts potent  
hypothermic and analgesic effects in the brain. In the periphery, neurotensin is a  
paracrine and endocrine modulator of the digestive tract and acts as a growth factor  
on a variety of cells.

So far, three types of neurotensin receptors have been identified:  
30 neurotensin-1 receptors, neurotensin-2 receptors, and neurotensin-3 receptors.  
(The neurotensin-3 receptor is also called sortlin or gp95.) The neurotensin-1 and  
neurotensin-2 receptors are G protein coupled receptors; the neurotensin-3 receptor  
is not a G protein coupled receptor.

### Summary of the Invention

The present invention provides methods of treating obesity, the methods comprising the step of administering to an obese patient or a patient at risk of becoming obese a therapeutically effective amount of a compound that is a neurotensin receptor ligand.

In a preferred embodiment of the methods, the neurotensin receptor ligand is a neurotensin-1 receptor ligand.

In another preferred embodiment of the methods, the neurotensin receptor ligand is a neurotensin-2 receptor ligand.

In another preferred embodiment of the methods, the neurotensin receptor ligand is a neurotensin-3 receptor ligand.

In another preferred embodiment of the methods, the ligand is an agonist.

In another preferred embodiment of the methods, the ligand is an antagonist.

In another preferred embodiment of the methods, the neurotensin receptor ligand is a neurotensin-1 receptor agonist.

Also provided are methods of treating obesity, the methods comprising the step of administering to an obese patient or a patient at risk of becoming obese a therapeutically effective amount of a compound that is a selective neurotensin-1 receptor agonist.

Also provided are pharmaceutical compositions comprising:

- a) a compound that is a neurotensin receptor ligand; and
- b) a second compound useful for the treatment of obesity, diabetes, sexual dysfunction, atherosclerosis, insulin resistance, impaired glucose tolerance, hypercholesterolemia or hypertriglyceridemia.

In a preferred embodiment of the compositions, the neurotensin receptor ligand is a neurotensin-1 receptor agonist.

In another preferred embodiment of the method, the second compound is a  $\beta_3$ -adrenergic receptor agonist, a cholecystokinin-A agonist, a monoamine reuptake inhibitor, a sympathomimetic agent, a serotonergic agent, a dopamine agonist, a melanocyte-stimulating hormone receptor agonist or mimetic, a melanocyte-stimulating hormone receptor analog, a cannabinoid receptor antagonist, a melanin concentrating hormone antagonist, leptin, a leptin analog, a leptin receptor agonist, a galanin antagonist, a bombesin agonist, a neuropeptide-Y antagonist, a thyromimetic agent, dehydroepiandrosterone or an analog thereof, a glucocorticoid receptor

agonist or antagonist, an orexin receptor antagonist, a urocortin binding protein antagonist, a glucagon-like peptide-1 receptor agonist, or a ciliary neurotrophic factor.

Also provided are kits that comprises:

- a) a first pharmaceutical composition comprising a compound that is a neurotensin receptor ligand;
- b) a second pharmaceutical composition comprising a compound that is useful for the treatment of obesity, diabetes, sexual dysfunction, atherosclerosis, insulin resistance, impaired glucose tolerance, hypercholesterolemia or hypertriglyceridemia; and
- c) a container for the first and second compositions.

In a preferred embodiment of the kits, the neurotensin receptor ligand is a neurotensin-1 receptor agonist.

In another preferred embodiment of the kits, the second pharmaceutical composition comprises a compound that is a  $\beta_3$ -adrenergic receptor agonist, a cholecystikinin-A agonist, a monoamine reuptake inhibitor, a sympathomimetic agent, a serotonergic agent, a dopamine agonist, a melanocyte-stimulating hormone receptor agonist or mimetic, a melanocyte-stimulating hormone receptor analog, a cannabinoid receptor antagonist, a melanin concentrating hormone antagonist, leptin, a leptin analog, a leptin receptor agonist, a galanin antagonist, a bombesin agonist, a neuropeptide-Y antagonist, a thyromimetic agent, dehydroepiandrosterone or an analog thereof, a glucocorticoid receptor agonist or antagonist, an orexin receptor antagonist, a urocortin binding protein antagonist, a glucagon-like peptide-1 receptor agonist, or a ciliary neurotrophic factor.

Also provided are methods of treating diabetes, sexual dysfunction, atherosclerosis, insulin resistance, impaired glucose tolerance, hypercholesterolemia or hypertriglyceridemia, the methods comprising the step of administering to a patient having or at risk of having, diabetes, sexual dysfunction, atherosclerosis, insulin resistance, impaired glucose tolerance, hypercholesterolemia or hypertriglyceridemia a therapeutically effective amount of a neurotensin receptor ligand.

In a preferred embodiment of the method, the neurotensin receptor ligand is a neurotension-1 receptor ligand.

#### Detailed Description of the Invention

The present invention relates to methods of treating obesity, diabetes, sexual dysfunction (including erectile dysfunction), atherosclerosis, insulin resistance,

impaired glucose tolerance, hypercholesterolemia or hypertriglyceridemia using a compound that is a neurotensin receptor ligand. In addition, the present invention provides pharmaceutical compositions and kits comprising a neurotensin receptor ligand.

5 In accordance with the present invention, obesity, diabetes, sexual dysfunction (including erectile dysfunction), atherosclerosis, insulin resistance, impaired glucose tolerance, hypercholesterolemia or hypertriglyceridemia can be treated by administering to an obese patient or a patient at risk of becoming obese or a patient having or at risk of having, diabetes, sexual dysfunction (including erectile  
10 dysfunction), atherosclerosis, insulin resistance, impaired glucose tolerance, hypercholesterolemia, or hypertriglyceridemia a therapeutically effective amount of a neurotensin receptor ligand. In a preferred embodiment of the invention, the neurotensin receptor ligand is a neurotensin-1 receptor ligand. In a more preferred embodiment of the invention, the neurotensin receptor ligand is a selective  
15 neurotensin-1 receptor agonist.

The term "therapeutically effective amount" means an amount of a compound or combination of compounds that treats a disease; ameliorates, attenuates, or eliminates one or more symptoms of a particular disease; or prevents or delays the onset of one or more symptoms of a disease.

20 The term "patient" means animals, such as dogs, cats, cows, horses, sheep, geese, and humans. Particularly preferred patients are mammals, including humans of both sexes.

The term "pharmaceutically acceptable" means that the substance or composition must be compatible with the other ingredients of a formulation, and not  
25 deleterious to the patient.

The terms "treating", "treat" or "treatment" include preventative (e.g., prophylactic) and palliative treatment.

The phrase "neurotensin receptor ligand" means a compound that binds to a neurotensin receptor, or a stereoisomer of the compound, a pharmaceutically  
30 acceptable salt of the compound or stereoisomer, a prodrug of the compound or stereoisomer, or a pharmaceutically acceptable salt of the prodrug. It is also contemplated that any additional pharmaceutically active compound used in combination with a neurotensin receptor ligand can be a stereoisomer of the additional active compound, a salt of the additional active compound or stereoisomer

thereof, a prodrug of the additional compound or stereoisomer thereof, or a salt of the prodrug.

The phrase "neurotensin receptor agonist" means a neurotensin receptor ligand that activates a neurotensin receptor.

5       The phrase "neurotensin receptor antagonist" means a neurotensin receptor ligand that blocks activation of a neurotensin receptor.

10       The term "selective" means that a ligand binds with greater affinity to a particular receptor when compared with the binding affinity of the ligand to another receptor. Preferably, the binding affinity of the ligand for the first receptor is about 50% or greater than the binding affinity for the second receptor. More preferably, the binding affinity of the ligand to the first receptor is about 75% or greater than the binding affinity to the second receptor. Most preferably, the binding affinity of the ligand to the first receptor is about 90% or greater than the binding affinity to the second receptor. In a preferred embodiment of the invention, the ligand exhibits a  
15       greater binding affinity to one of the three neurotensin receptors. Particularly preferred ligands are those that bind with greater affinity to the neurotensin-1 receptors when compared with binding to the neurotensin-2 or neurotensin-3 receptors. It is contemplated that preferred compounds bind neurotensin receptors with micromolar or greater affinity. More preferred compounds bind neurotensin  
20       receptors with nanomolar or greater affinity. Preferred neurotensin receptor ligands of the present invention include compounds that are selective agonists of the neurotensin-1 receptor.

25       Neurotensin receptor ligands can be identified, for example, by screening a compound library. Methods of identifying agonists and antagonists of receptors are well known to those skilled in the art. Specific procedures that can be used to identify neurotensin receptor ligands are presented below.

30       Examples of known neurotensin receptor ligands include hormones such as neurotensin (also called NT(1-13)) and neuromedin N, non-peptide agonists such as those disclosed in U.S. patent number 5,407,916, non-peptide antagonists such as 2-  
3yl]carboxylamino)tricyclo(3.3.1.1.[3.7)  
decan-2-carboxylic acid (SR48692), which is a selective neurotensin-1 receptor antagonist, and 2-(5,6-dimethylaminopropyl)-1-[4-{N-(3-dimethylaminopropyl)-N-methylcarbamoyl}-2-isopropylphenyl]-1H-pyrazole-3-carbonyl)aminoadamantane-2-

carboxylic acid (SR142948A), which is a non-selective antagonist that binds with equal affinity at neurotensin-1 and neurotensin-2 receptors. Another compound that is a neurotensin binding ligand is levocabastine. In addition, U.S. patents 5,250,558 and 5,204,354 disclose neurotensin receptor antagonists, and U.S. patent 5,407,916 discloses peptidic neurotensin agonists. An example of a selective neurotensin-1 receptor agonist is native neurotensin [NT(1-13)], which has a  $K_d$  of about 0.3 nM at the neurotensin-1 receptor and about 2-6 nM at the neurotensin-2 receptor. Another example of a selective neurotensin-1 receptor agonist is Trp11 NT(1-13), which shows a binding affinity of about 1 nM at the neurotensin-1 receptor and about 27 nM at the neurotensin-2 receptor. Trp11 NT(1-13) is NT(1-13) in which amino acid 11 is tryptophan.

The amino acid sequences and nucleotide sequences that encode each of the three human neurotensin receptors are known to those skilled in the art and can be found in GenBank under accession numbers NM\_002531, Y10148, and NM\_002569.

A neurotensin receptor ligand is administered to a patient in a therapeutically effective amount. A neurotensin receptor ligand can be administered alone or as part of a pharmaceutically acceptable composition. In addition, a compound or composition can be administered all at once, as for example, by a bolus injection, multiple times, such as by a series of tablets, or delivered substantially uniformly over a period of time, as for example, using transdermal delivery. It is also noted that the dose of the compound can be varied over time. A neurotensin receptor ligand can be administered using an immediate release formulation, a controlled release formulation, or combinations thereof. The term "controlled release" includes sustained release, delayed release, and combinations thereof.

In addition, a neurotensin receptor ligand can be administered alone, in combination with other neurotensin receptor ligands, or with other pharmaceutically active compounds. The other pharmaceutically active compounds can be intended to treat the same disease as the neurotensin receptor ligand or a different disease. If the patient is to receive or is receiving multiple pharmaceutically active compounds, the compounds can be administered simultaneously or sequentially in any order. For example, in the case of tablets, the active compounds may be found in one tablet or in separate tablets, which can be administered at once or sequentially in any order. In addition, it should be recognized that the compositions can be different forms. For

example, one or more compounds may be delivered via a tablet, while another is administered via injection or orally as a syrup. All combinations, delivery methods and administration sequences are contemplated.

Since one aspect of the present invention contemplates the treatment of the  
5 diseases referenced with a combination of pharmaceutically active agents that may be administered separately, the invention further relates to combining separate pharmaceutical compositions in kit form. For example, a kit may comprise two separate pharmaceutical compositions comprising: 1) a neurotensin receptor ligand; and 2) a second pharmaceutically active compound. The kit also comprises a  
10 container for the separate compositions, such as a divided bottle or a divided foil packet. Additional examples of containers include syringes, boxes, bags, and the like. Typically, a kit comprises directions for the administration of the separate components. The kit form is particularly advantageous when the separate components are preferably administered in different dosage forms (e.g., oral and parenteral), are administered at different dosage intervals, or when titration of the  
15 individual components of the combination is desired by the prescribing physician.

An example of a kit is a blister pack. Blister packs are well known in the packaging industry and are being widely used for the packaging of pharmaceutical unit dosage forms (tablets, capsules, and the like). Blister packs generally consist of  
20 a sheet of relatively stiff material covered with a foil of a preferably transparent plastic material. During the packaging process recesses are formed in the plastic foil. The recesses have the size and shape of the tablets or capsules to be packed. Next, the tablets or capsules are placed in the recesses and a sheet of relatively stiff material is sealed against the plastic foil at the face of the foil which is opposite from the  
25 direction in which the recesses were formed. As a result, the tablets or capsules are sealed in the recesses between the plastic foil and the sheet. Preferably the strength of the sheet is such that the tablets or capsules can be removed from the blister pack by manually applying pressure on the recesses whereby an opening is formed in the sheet at the place of the recess. The tablet or capsule can then be removed via said  
30 opening.

It may be desirable to provide a memory aid on the kit, e.g., in the form of numbers next to the tablets or capsules whereby the numbers correspond with the days of the regimen that the tablets or capsules so specified should be ingested. Another example of such a memory aid is a calendar printed on the card, e.g., as

follows "First Week, Monday, Tuesday, ...etc.... Second Week, Monday, Tuesday," etc. Other variations of memory aids will be readily apparent. A "daily dose" can be a single tablet or capsule or several pills or capsules to be taken on a given day. Also, a daily dose of a neurotensin receptor ligand can consist of one tablet or capsule, while a daily dose of the second compound can consist of several tablets or capsules and vice versa. The memory aid should reflect this and assist in correct administration of the active agents.

In another embodiment of the present invention, a dispenser designed to dispense the daily doses one at a time in the order of their intended use is provided. Preferably, the dispenser is equipped with a memory aid, so as to further facilitate compliance with the dosage regimen. An example of such a memory aid is a mechanical counter, which indicates the number of daily doses that have been dispensed. Another example of such a memory aid is a battery-powered micro-chip memory coupled with a liquid crystal readout, or audible reminder signal which, for example, reads out the date that the last daily dose has been taken and/or reminds one when the next dose is to be taken.

A neurotensin receptor ligand and other pharmaceutically active compounds, if desired, can be administered to a patient either orally, rectally, parenterally, (for example, intravenously, intramuscularly, or subcutaneously) intracisternally, intravaginally, intraperitoneally, intravesically, locally (for example, powders, ointments or drops), or as a buccal or nasal spray.

Compositions suitable for parenteral injection may comprise physiologically acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions, or emulsions, or may comprise sterile powders for reconstitution into sterile injectable solutions or dispersions. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents, or vehicles include water, ethanol, polyols (propylene glycol, polyethylene glycol, glycerol, and the like), suitable mixtures thereof, triglycerides, including vegetable oils such as olive oil, or injectable organic esters such as ethyl oleate. A preferred carrier is Miglyol®. Proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersions, and/or by the use of surfactants.

These compositions may also contain adjuvants such as preserving, wetting, emulsifying, and/or dispersing agents. Prevention of microorganism contamination of the compositions can be accomplished by the addition of various antibacterial and

Patent 3,443,660



antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, and the like. It may also be desirable to include isotonic agents, for example, sugars, sodium chloride, and the like. Prolonged absorption of injectable pharmaceutical compositions can be brought about by the use of agents capable of delaying absorption, for example, aluminum monostearate and/or gelatin.

5 Solid dosage forms for oral administration include capsules, tablets, powders, and granules. In such solid dosage forms, the active compound is admixed with at least one inert customary excipient (or carrier) such as sodium citrate or dicalcium phosphate or (a) fillers or extenders, as for example, starches, lactose, sucrose, 10 mannitol, or silicic acid; (b) binders, as for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose, or acacia; (c) humectants, as for example, glycerol; (d) disintegrating agents, as for example, agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain complex silicates, or sodium carbonate; (e) solution retarders, as for example, paraffin; (f) absorption accelerators, 15 as for example, quaternary ammonium compounds; (g) wetting agents, as for example, cetyl alcohol or glycerol monostearate; (h) adsorbents, as for example, kaolin or bentonite; and/or (i) lubricants, as for example, talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, or mixtures thereof. In the case of capsules and tablets, the dosage forms may also comprise 20 buffering agents.

Solid compositions of a similar type may also be used as fillers in soft or hard filled gelatin capsules using such excipients as lactose or milk sugar, as well as high molecular weight polyethylene glycols, and the like.

25 Solid dosage forms such as tablets, dragees, capsules, and granules can be prepared with coatings or shells, such as enteric coatings and others well known in the art. They may also contain opacifying agents, and can also be of such composition that they release the active compound or compounds in a delayed manner. Examples of embedding compositions that can be used are polymeric substances and waxes. The active compounds can also be in micro-encapsulated 30 form, if appropriate, with one or more of the above-mentioned excipients.

Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs. In addition to the active compounds, the liquid dosage form may contain inert diluents commonly used in the art, such as water or other solvents, solubilizing agents and emulsifiers, as for

example, ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils, in particular, cottonseed oil, groundnut oil, corn germ oil, olive oil, castor oil, sesame seed oil, Miglyol<sup>®</sup>, glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols, fatty acid esters of sorbitan, or mixtures of these substances, and the like.

Besides such inert diluents, the composition can also include adjuvants, such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

Suspensions, in addition to the active compound, may contain suspending agents, as for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol or sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar, or tragacanth, or mixtures of these substances, and the like.

Compositions for rectal or vaginal administration can be prepared by mixing a neurotensin receptor ligand and any additional compounds with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax, which are solid at ordinary room temperature, but liquid at body temperature, and therefore, melt in the rectum or vaginal cavity and release the neurotensin receptor ligand.

Dosage forms for topical administration of a neurotensin receptor ligand include ointments, powders, sprays and inhalants. The compound(s) are admixed under sterile conditions with a physiologically acceptable carrier, and any preservatives, buffers, and/or propellants that may be required. Ophthalmic formulations, eye ointments, powders, and solutions are also contemplated as being within the scope of this invention.

A neurotensin receptor ligand can be administered to a patient at dosage levels in the range of about 0.1 to about 7,000 mg per day. A preferred dosage range is about 1 to about 100 mg per day. The specific dosage and dosage range that can be used depends on a number of factors, including the requirements of the patient, the severity of the condition or disease being treated, and the pharmacological activity of the compound being administered. The determination of dosage ranges and optimal dosages for a particular patient is well within the ordinary skill of one in the art in view of this disclosure.

The following paragraphs describe exemplary formulations, dosages,

An amount of a neurotensin receptor ligand is administered such that a therapeutically effective dose is received, generally a daily dose which, when

Conveniently, the compound or compounds can be carried in the drinking water so that a therapeutic dose of the compound or compounds is ingested

Conveniently, the compound or compounds can also be added directly to the feed, as such, or in the form of an animal feed supplement, also referred to as

15 commonly employed for the inclusion of the compound or compounds in the feed. Suitable carriers are liquid or solid, as desired, such as water, various

mixes such as are commonly employed in poultry feeds. A particularly  
20 effective carrier is the respective animal feed itself; that is, a small portion of

important that the compound or compounds be thoroughly blended into the premix and, subsequently, the feed. In this respect, the compound or

solvent and then blended with the carrier. It will be appreciated that the proportions of compound or compounds in the concentrate are capable of

High potency concentrates may be blended by the feed manufacturer with proteinaceous carrier such as soybean oil meal or other meals, as described above, to produce concentrated supplements that are suitable for direct feeding to animals.

High potency concentrates may be blended by the feed manufacturer with proteinaceous carrier such as soybean oil meal or other meals, as described above, to produce concentrated supplements that are suitable for direct feeding to animals.

5

d

10

in

15

20

25

30

include cations based on the alkali and alkaline earth metals, such as sodium, lithium, potassium, calcium, magnesium, and the like, as well as non-toxic ammonium, quaternary ammonium, and amine cations including, but not limited to, ammonium, tetramethylammonium, tetraethylammonium, methylamine, dimethylamine, trimethylamine, triethylamine, ethylamine, and the like. See, for example, S.M. Berge, et al., "Pharmaceutical Salts," *J Pharm Sci*, **66**:1-19 (1977).

The term "prodrug" means a compound that is transformed *in vivo* to yield a therapeutically active compound. The transformation may occur by various mechanisms, such as through hydrolysis in blood. A discussion of the use of prodrugs is provided by T. Higuchi and W. Stella, "Pro-drugs as Novel Delivery Systems," Vol. 14 of the A.C.S. *Symposium Series*, and in *Bioreversible Carriers in Drug Design*, ed. Edward B. Roche, American Pharmaceutical Association and Pergamon Press, 1987.

For example, if a compound contains a carboxylic acid functional group, a prodrug can comprise an ester formed by the replacement of the hydrogen atom of the acid group with a group such as (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>2</sub>-C<sub>12</sub>)alkanoyloxymethyl, 1-(alkanoyloxy)ethyl having from 4 to 9 carbon atoms, 1-methyl-1-(alkanoyloxy)-ethyl having from 5 to 10 carbon atoms, alkoxycarbonyloxymethyl having from 3 to 6 carbon atoms, 1-(alkoxycarbonyloxy)ethyl having from 4 to 7 carbon atoms, 1-methyl-1-(alkoxycarbonyloxy)ethyl having from 5 to 8 carbon atoms, N-(alkoxycarbonyl)aminomethyl having from 3 to 9 carbon atoms, 1-(N-(alkoxycarbonyl)amino)ethyl having from 4 to 10 carbon atoms, 3-phthalidyl, 4-crotonolactonyl, gamma-butyrolacton-4-yl, di-N,N-(C<sub>1</sub>-C<sub>2</sub>)alkylamino(C<sub>2</sub>-C<sub>3</sub>)alkyl (such as β-dimethylaminoethyl), carbamoyl-(C<sub>1</sub>-C<sub>2</sub>)alkyl, N,N-di(C<sub>1</sub>-C<sub>2</sub>)alkylcarbamoyl-(C<sub>1</sub>-C<sub>2</sub>)alkyl and piperidino-, pyrrolidino- or morpholino(C<sub>2</sub>-C<sub>3</sub>)alkyl.

Similarly, if a compound comprises an alcohol functional group, a prodrug can be formed by the replacement of the hydrogen atom of the alcohol group with a group such as (C<sub>1</sub>-C<sub>6</sub>)alkanoyloxymethyl, 1-((C<sub>1</sub>-C<sub>6</sub>)alkanoyloxy)ethyl, 1-methyl-1-((C<sub>1</sub>-C<sub>6</sub>)alkanoyloxy)ethyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyloxymethyl, N-(C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylaminomethyl, succinoyl, (C<sub>1</sub>-C<sub>6</sub>)alkanoyl, α-amino(C<sub>1</sub>-C<sub>4</sub>)alkanoyl, arylacyl and α-aminoacyl, or α-aminoacyl-α-aminoacyl, where each α-aminoacyl group is independently selected from the naturally occurring L-amino acids, P(O)(OH)<sub>2</sub>, -P(O)(O(C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub> or glycosyl (the radical resulting from the removal of a hydroxyl group of the hemiacetal form of a carbohydrate).

If a compound comprises an amine functional group, a prodrug can be formed by the replacement of a hydrogen atom in the amine group with a group such as R-carbonyl, RO-carbonyl, NRR'-carbonyl where R and R' are each independently ((C<sub>1</sub>-C<sub>10</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, benzyl, or R-carbonyl is a natural  $\alpha$ -aminoacyl or natural  $\alpha$ -aminoacyl-natural  $\alpha$ -aminoacyl, -C(OH)C(O)OY wherein Y is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl or benzyl, -C(OY<sub>0</sub>)Y<sub>1</sub> wherein Y<sub>0</sub> is (C<sub>1</sub>-C<sub>4</sub>) alkyl and Y<sub>1</sub> is ((C<sub>1</sub>-C<sub>6</sub>)alkyl, carboxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>4</sub>)alkyl or mono-N- or di-N,N-(C<sub>1</sub>-C<sub>6</sub>)alkylaminoalkyl, -C(Y<sub>2</sub>)Y<sub>3</sub> wherein Y<sub>2</sub> is H or methyl and Y<sub>3</sub> is mono-N- or di-N,N-(C<sub>1</sub>-C<sub>6</sub>)alkylamino, morpholino, piperidin-1-yl or pyrrolidin-1-yl.

10 A neurotensin receptor ligand may contain asymmetric or chiral centers, and therefore, exist in different stereoisomeric forms. It is contemplated that all stereoisomeric forms as well as mixtures thereof, including racemic mixtures, form part of the present invention. In addition, the present invention contemplates all geometric and positional isomers. For example, if a compound contains a double  
15 bond, both the cis and trans forms, as well as mixtures, are contemplated.

Mixtures of isomers, including stereoisomers can be separated into their individual isomers on the basis of their physical chemical differences by methods well know to those skilled in the art, such as by chromatography and/or fractional crystallization. Enantiomers can be separated by converting the enantiomeric mixture  
20 into a diastereomeric mixture by reaction with an appropriate optically active compound (e.g., alcohol), separating the diastereomers and converting (e.g., hydrolyzing) the individual diastereomers to the corresponding pure enantiomers. Also, some of the compounds of this invention may be atropisomers (e.g., substituted biaryls) and are considered as part of this invention.

25 A neurotensin receptor ligand may exist in unsolvated as well as solvated forms with pharmaceutically acceptable solvents such as water, ethanol, and the like. The present invention contemplates and encompasses both the solvated and unsolvated forms.

It is also possible that a neurotensin receptor ligand may exist in different  
30 tautomeric forms. All tautomers of a neurotensin receptor ligand are contemplated.

Those skilled in the art will recognize that compound names contained herein may be based on a particular tautomer of a compound. While the name for only a particular tautomer may be used, it is intended that all tautomers are encompassed

by the name of the particular tautomer, and all tautomers are considered part of the present invention.

It is also intended that the invention disclosed herein encompass compounds that are synthesized *in vitro* using laboratory techniques, such as those well known to synthetic chemists; or synthesized using *in vivo* techniques, such as through metabolism, fermentation, digestion, and the like. It is also contemplated that compounds may be synthesized using a combination of *in vitro* and *in vivo* techniques.

The present invention also includes isotopically labeled compounds, which are identical to the non-isotopically labeled compounds, but for the fact that one or more atoms are replaced by an atom having an atomic mass or mass number different from the atomic mass or mass number usually found most abundantly in nature. Examples of isotopes that can be incorporated into compounds identified by the present invention include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, fluorine and chlorine, such as  $^2\text{H}$ ,  $^3\text{H}$ ,  $^{13}\text{C}$ ,  $^{14}\text{C}$ ,  $^{15}\text{N}$ ,  $^{18}\text{O}$ ,  $^{17}\text{O}$ ,  $^{31}\text{P}$ ,  $^{32}\text{P}$ ,  $^{35}\text{S}$ ,  $^{18}\text{F}$ ,  $^{135}\text{I}$  and  $^{36}\text{Cl}$ , respectively. Neurotensin receptor ligands, prodrugs thereof, and pharmaceutically acceptable salts of said ligands or of said prodrugs which contain the aforementioned isotopes and/or other isotopes of other atoms are within the scope of this invention. Certain isotopically labeled compounds of the present invention, for example those into which radioactive isotopes such as  $^3\text{H}$  and  $^{14}\text{C}$  are incorporated, are useful in drug and/or substrate tissue distribution assays. Tritiated, i.e.,  $^3\text{H}$ , and carbon-14, i.e.,  $^{14}\text{C}$ , isotopes are particularly preferred for their ease of preparation and detectability. Further, substitution with heavier isotopes such as deuterium, i.e.,  $^2\text{H}$ , may afford certain therapeutic advantages resulting from greater metabolic stability, for example increased *in vivo* half-life or reduced dosage requirements and, hence, may be preferred in some circumstances. Isotopically labeled compounds can generally be prepared by substituting a readily available isotopically labeled reagent for a non-isotopically labeled reagent.

The present invention relates to the use of neurotensin receptor ligands to treat obesity, diabetes, sexual dysfunction, atherosclerosis, insulin resistance, impaired glucose tolerance, hypercholesterolemia, or hypertriglyceridemia.

In addition, a neurotensin receptor ligand, particularly neurotensin-2 receptor ligands, can be used to treat hyperthermia; hypothermia; gastrointestinal ulcers; substance abuse; depression; Alzheimer's disease; tardive dyskinesia; panic attack;

gastrointestinal reflux disorder; irritable bowel syndrome; diarrhea; cholic; dyspepsia; pancreatitis; esophagitis; gastroparesis; neurological diseases such as schizophrenia, psychoses, anxiety, manic depression, delirium dementia, severe mental retardation, and dyskinesias such as Huntington's disease and Tourett's syndrome; fungal and viral infections, including HIV-1 and HIV-2 infections; pain (i.e., an analgesic); cancer (including gastrointesitinal tumors); anorexia; bulimia; asthma; Parkinson's disease; acute heart failure; hypotension; hypertension; urinary retention; osteoporosis; angina pectoris; myocardial infarction; allergies; inflammation; or benign prostatic hypertrophy.

10       The methods of treatment of the present invention can also include combination therapy where other pharmaceutically active compounds useful for the treatment of obesity or other diseases are used in combination with a neurotensin receptor ligand.

15       It is known that obese patients have higher incidences of certain diseases such as atherosclerosis, hypercholesterolemia, hypertriglyceridemia, hypertension, sexual dysfunction (including erectile dysfunction), insulin resistance, impaired glucose tolerance, diabetes, [particularly non-insulin dependent diabetes mellitus (NIDDM or Type 2 diabetes)] and the diseases associated with diabetes such as nephropathy, neuropathy, retinopathy, cardiomyopathy, cataracts, and polycystic  
20       ovary syndrome. These diseases can be treated indirectly by treating obesity using a neurotensin receptor ligand or directly by treating the specific disease itself using a neurotensin receptor ligand. These diseases can be treated in the absence of obesity using a neurotensin receptor ligand.

25       In one embodiment of the invention, an obese patient or a patient at risk of becoming obese can be administered a combination of: 1) a neurotensin receptor ligand; and 2) an additional compound useful to treat obesity, diabetes [including (NIDDM) and the conditions and/or diseases associated with diabetes, such as nephropathy, neuropathy, retinopathy, cardiomyopathy, cataracts, and polycystic ovary syndrome], atherosclerosis, hypercholesterolemia, hypertriglyceridemia, sexual  
30       dysfunction (including erectile dysfunction), insulin resistance, or impaired glucose tolerance, or combinations of compounds useful to treat these diseases.

Sexual dysfunction occurs in males and females and includes hypoactive sexual desire disorder, sexual anhedonia and dyspareunia. Hypoactive sexual desire disorder is a disorder in which sexual fantasies and desire for sexual activity are

the "obese"



persistently or recurrently diminished or absent, causing marked distress or interpersonal difficulties. Symptoms and signs of hypoactive sexual desire disorder include the patient complaining of a lack of interest in sex, even in ordinarily erotic situations. The disorder is usually associated with infrequent sexual activity, often causing serious conflict between partners. Sexual anhedonia is decreased or absent pleasure in sexual activity. Sexual anhedonia is almost always classified under hypoactive sexual desire disorder, because loss of pleasure typically results in loss of desire. Dyspareunia is painful coitus or attempted coitus.

Erectile dysfunction is another example of a sexual dysfunction. Erectile dysfunction, like obesity, is another condition that can result in severe emotional distress. Persons suffering from erectile dysfunction are unable to develop and/or maintain an erection of the penis. Historically, erectile dysfunction has been viewed as having biological and psychological components, and more effort appeared to be exerted on treating the psychological components of the condition. Only recently with the introduction of Viagra® have persons having this condition been offered an oral medicinal treatment.

Diabetes is found more frequently in obese patients than non-obese patients.

In spite of the early discovery of insulin and its subsequent widespread use in the treatment of diabetes, and the later discovery of and use of sulfonylureas, biguanides and thiazolidenediones, such as troglitazone, rosiglitazone or pioglitazone, as oral hypoglycemic agents, the treatment of diabetes remains less than satisfactory.

The use of insulin currently requires multiple daily doses, usually by self-injection. Determination of the proper dosage of insulin requires frequent estimations of the sugar in urine or blood. The administration of an excess dose of insulin causes hypoglycemia, with effects ranging from mild abnormalities in blood glucose to coma, or even death. Treatment of non-insulin dependent diabetes mellitus (Type 2 diabetes, NIDDM) usually consists of a combination of diet, exercise, oral hypoglycemic agents, e.g., thiazolidenediones, and, in more severe cases, insulin. However, the clinically available hypoglycemic agents can have side effects that limit their use, or an agent may not be effective with a particular patient. In the case of insulin dependent diabetes mellitus (Type 1), insulin is usually the primary course of therapy. Additional hypoglycemic agents that have fewer side effects or succeed where others fail are needed.

Atherosclerosis, a disease of the arteries, is recognized to be a leading cause of death in the United States and Western Europe. The pathological sequence leading to atherosclerosis and occlusive heart disease is well known. The earliest stage in this sequence is the formation of "fatty streaks" in the carotid, coronary and cerebral arteries and in the aorta. These lesions are yellow in color due to the presence of lipid deposits found principally within smooth-muscle cells and in macrophages of the intima layer of the arteries and aorta. Further, it is postulated that most of the cholesterol found within the fatty streaks, in turn, give rise to development of "fibrous plaques," which consist of accumulated intimal smooth muscle cells laden with lipid and are surrounded by extra-cellular lipid, collagen, elastin and proteoglycans. The cells plus matrix form a fibrous cap that covers a deeper deposit of cell debris and more extra-cellular lipid. The lipid is primarily free and esterified cholesterol. The fibrous plaque forms slowly, and is likely in time to become calcified and necrotic, advancing to a "complicated lesion," which accounts for arterial occlusion and tendency toward mural thrombosis and arterial muscle spasm that characterize advanced atherosclerosis.

Epidemiological evidence has firmly established hyperlipidemia as a primary risk factor in causing cardiovascular disease (CVD) due to atherosclerosis. In recent years, leaders of the medical profession have placed renewed emphasis on lowering plasma cholesterol levels, and low density lipoprotein cholesterol in particular, as an essential step in prevention of CVD. The upper limits of "normal" are now known to be significantly lower than heretofore appreciated. As a result, large segments of Western populations are now realized to be at particularly high risk. Such independent risk factors include glucose intolerance, left ventricular hypertrophy, hypertension, and being of the male sex. Cardiovascular disease is especially prevalent among diabetic subjects, at least in part because of the existence of multiple independent risk factors in this population. Successful treatment of hyperlipidemia in the general population, and in diabetic subjects in particular, is therefore of exceptional medical importance.

Hypertension (or high blood pressure) is a condition that occurs in the human population as a secondary symptom to various other disorders such as renal artery stenosis, pheochromocytoma or endocrine disorders. However, hypertension is also evidenced in many patients in whom the causative agent or disorder is unknown. While such "essential" hypertension is often associated with disorders such as

104-10860-10

It is known that hypertension can directly lead to heart failure, renal failure and stroke (brain hemorrhaging). These conditions are capable of causing death in a patient. Hypertension can also contribute to the development of atherosclerosis and coronary disease. These conditions gradually weaken a patient and can lead to death.

15           The treatment of essential hypertension has been undertaken bearing the foregoing factors in mind. Thus, a broad range of beta-blockers, vasoconstrictors, angiotensin converting enzyme inhibitors and the like have been developed and marketed as antihypertensives. The treatment of hypertension utilizing these compounds has proven beneficial in the prevention of short-interval deaths such as

20   heart failure, renal failure and brain hemorrhaging.

A neurotensin receptor ligand can be used in combination with one or more compounds that are useful to treat obesity. Examples of classes of compounds that can be used to treat obesity include the active compound(s) in appetite suppressants

A neurotensin receptor ligand can be used in combination with one or more compounds that are useful to treat obesity. Examples of classes of compounds that can be used to treat obesity include the active compound(s) in appetite suppressants

such as Adipex®, Bontril®, Desoxyn Gradumet®, Fastin®, Ionamin®, and Meridia®, and lipase inhibitors such as Xenical®.

Additional anti-obesity agents that can be used in combination with a neurotensin receptor ligand include a  $\beta_3$ -adrenergic receptor agonist, a cholecystokinin-A agonist, a monoamine reuptake inhibitor, a sympathomimetic agent, a serotonergic agent, a dopamine agonist, a melanocyte-stimulating hormone receptor agonist or mimetic, a melanocyte-stimulating hormone receptor analog, a cannabinoid receptor antagonist, a melanin concentrating hormone antagonist, leptin, a leptin analog, a leptin receptor agonist, a galanin antagonist, a bombesin agonist, a neuropeptide-Y antagonist (including NPY-1 and NPY-5), a thyromimetic agent, dehydroepiandrosterone or an analog thereof, a glucocorticoid receptor agonist or antagonist, an orexin receptor antagonist, a urocortin binding protein antagonist, a glucagon-like peptide-1 receptor agonist, and a ciliary neurotrophic factor.

Especially preferred anti-obesity agents that can be used in combination with a neurotensin receptor ligand include compounds selected from the group consisting of sibutramine, fenfluramine, dexfenfluramine, bromocriptine, phentermine, orlistat, ephedrine, leptin, phenylpropanolamine, pseudoephedrine, {4-[2-(2-[6-aminopyridin-3-yl]-2(R)-hydroxyethylamino)ethoxy]phenyl}acetic acid, {4-[2-(2-[6-aminopyridin-3-yl]-2(R)-hydroxyethylamino)ethoxy]phenyl}benzoic acid, {4-[2-(2-[6-aminopyridin-3-yl]-2(R)-hydroxyethylamino)ethoxy]phenyl}propionic acid, and {4-[2-(2-[6-aminopyridin-3-yl]-2(R)-hydroxyethylamino)ethoxy]phenoxy}acetic acid.

Examples of thyromimetics that can be used in combination with a neurotensin receptor ligand include those disclosed in U.S. provisional patent application numbers 60/178,968 and 60/177,987.

Examples of glucocorticoid receptor ligands that can be used in combination with a neurotensin receptor ligand include those disclosed in U.S. provisional patent application number 60/132,130.

Examples of neuropeptide-Y antagonists that can be used in combination with a neurotensin receptor ligand include those disclosed in WO 98/23603, U.S. 5,900,415, U.S. 5,914,329, and U.S. provisional patent application number 60/132,029 (NPY-5).

Examples of  $\beta_3$ -adrenergic receptor agonists that can be used in combination with a neurotensin receptor ligand include those disclosed in WO 96/35671.

Additional compounds that can be used to treat obesity and that can be used in combination with a neurotensin receptor ligand include the compounds disclosed in WO 98/46243.

Similarly, compounds that can be used to treat sexual dysfunction, and particularly erectile dysfunction, such as Viagra® can also be used in combination with a neurotensin receptor ligand. Other compounds that can be used to treat sexual dysfunction, particularly erectile dysfunction, and that can be used in combination with a neurotensin receptor ligand include apomorphine and IC351 (ICOS). A class of compounds that are useful to treat sexual dysfunction, particularly  
5  
10  
erectile dysfunction, are phosphodiesterase V inhibitors. Examples of phosphodiesterase V inhibitors can be found in U.S. patent number 5,272,147.

In another aspect of the invention, a neurotensin receptor ligand can be administered in combination with a compound that is known to treat hypertension. Examples of classes of compounds that can be used to treat hypertension include  
15  
calcium blockers, ACE inhibitors, diuretics, angiotensin II receptor blockers,  $\beta$ -blockers, and  $\alpha$ -adrenergic blockers. In addition, combinations of compounds in the above-recited classes have been used to treat hypertension. Some examples of specific compounds that can be used in combination with neurotensin receptor ligands include quinapril; amlodipine, including the besylate salt; nifedipine;  
20  
doxazosin, including the mesylate salt; and prazosin, including the hydrochloride salt.

In another aspect, a neurotensin receptor ligand can be used in combination with compounds useful for the treatment of diabetes, including impaired glucose tolerance, insulin resistance, insulin dependent diabetes mellitus (Type 1) and non-insulin dependent diabetes mellitus (NIDDM or Type 2). Also intended to be  
25  
encompassed in the treatment of diabetes are the diabetic complications, such as neuropathy, nephropathy, retinopathy, cardiomyopathy or cataracts.

Representative agents that can be used to treat diabetes and which can be used in combination with a neurotensin receptor ligand include but are not limited to insulin and insulin analogs (e.g., LysPro insulin); GLP-1 (7-37)  
30  
(insulintropin) and GLP-1 (7-36)-NH<sub>2</sub>; sulfonylureas and analogs: chlorpropamide, glibenclamide, tolbutamide, tolazamide, acetohexamide, glypizide, glimepiride, repaglinide, meglitinide; biguanides: metformin, phenformin, buformin;  $\alpha$ 2-antagonists and imidazolines: midaglizole, isaglidole, deriglidole, idazoxan, efaroxan, fluparoxan; other insulin

- secretagogues: linogiride, A-4166; glitazones: ciglitazone, pioglitazone, englitazone, troglitazone, darglitazone, BRL49653; fatty acid oxidation inhibitors: clomoxir, etomoxir;  $\alpha$ -glucosidase inhibitors: acarbose, miglitol, emiglitate, voglibose, MDL-25,637, camiglibose, MDL-73,945;  $\beta$ -agonists:
- 5 BRL 35135, BRL 37344, Ro 16-8714, ICI D7114, CL 316,243; phosphodiesterase inhibitors: L-386,398; lipid-lowering agents: benfluorex; antiobesity agents: fenfluramine and orlistat; vanadate and vanadium complexes (e.g., Naglivan<sup>®</sup>) and peroxovanadium complexes; amylin antagonists; glucagon antagonists; gluconeogenesis inhibitors; somatostatin
- 10 agonists and antagonists; antilipolytic agents: nicotinic acid, acipimox, WAG 994; and glycogen phosphorylase inhibitors, such as those disclosed in WO 96/39385 and WO 96/39384. Also contemplated in combination with compounds of the present invention are pramlintide acetate (Symlin<sup>™</sup>) and nateglinide.
- 15 Preferred examples of glycogen phosphorylase inhibitors that can be used in the present invention in combination with a neurotensin receptor ligand include: 6H-thieno[2,3-b]pyrrole-5-carboxylic acid [(1S)-benzyl-3-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-(2R)-hydroxy-3-oxo-propyl]-amide;
- 2-bromo-6H-thieno[2,3-b]pyrrole-5-carboxylic acid [(1S)-benzyl-3-((3R,4S)-
- 20 dihydroxy-pyrrolidin-1-yl)-(2R)-hydroxy-3-oxo-propyl]-amide;
- 2-methyl-6H-thieno[2,3-b]pyrrole-5-carboxylic acid [(1S)-benzyl-3-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-(2R)-hydroxy-3-oxo-propyl]-amide;
- ( $\pm$ )-2-methyl-6H-thieno[2,3-b]pyrrole-5-carboxylic acid [1-benzyl-2-((3R,4S)-
- dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide;
- 25 2-bromo-6H-thieno[2,3-b]pyrrole-5-carboxylic acid [(1S)-benzyl-2-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide;
- 2-chloro-6H-thieno[2,3-b]pyrrole-5-carboxylic acid [(1S)-benzyl-3-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-(2R)-hydroxy-3-oxo-propyl]-amide;
- 2-chloro-6H-thieno[2,3-b]pyrrole-5-carboxylic acid [(1S)-benzyl-2-((3R,4S)-
- 30 dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide;
- 2,4-dichloro-6H-thieno[2,3-b]pyrrole-5-carboxylic acid [(1S)-benzyl-3-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-(2R)-hydroxy-3-oxo-propyl]-amide;

- (±)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid [1-benzyl-2-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide;
- 2-bromo-4H-thieno[3,2-b]pyrrole-5-carboxylic acid [(1S)-benzyl-3-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-(2R)-hydroxy-3-oxo-propyl]-amide;
- 5 4H-thieno[3,2-b]pyrrole-5-carboxylic acid [(1S)-benzyl-3-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-(2R)-hydroxy-3-oxo-propyl]-amide;
- (±)-2-bromo-4H-furo[3,2-b]pyrrole-5-carboxylic acid [1-benzyl-2-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide;
- 2-bromo-4H-furo[3,2-b]pyrrole-5-carboxylic acid [(1S)-benzyl-3-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-(2R)-hydroxy-3-oxo-propyl]-amide;
- 10 6H-thieno[2,3-b]pyrrole-5-carboxylic acid [(1S)-benzyl-2-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide;
- 2-bromo-4H-thieno[3,2-b]pyrrole-5-carboxylic acid [(1S)-benzyl-2-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide;
- 15 2-methyl-4H-thieno[3,2-b]pyrrole-5-carboxylic acid [(1S)-benzyl-2-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide;
- 2,4-dichloro-6H-thieno[2,3-b]pyrrole-5-carboxylic acid [(1S)-benzyl-2-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide;
- 2-cyano-6H-thieno[2,3-b]pyrrole-5-carboxylic acid [(1S)-benzyl-2-(3-hydroxy-azetidin-1-yl)-2-oxo-ethyl]-amide;
- 20 1-yl)-2-oxo-ethyl]-amide;
- 2-chloro-6H-thieno[2,3-b]pyrrole-5-carboxylic acid [(1S)-benzyl-2-morpholin-4-yl-2-oxo-ethyl]-amide;
- 2-chloro-6H-thieno[2,3-b]pyrrole-5-carboxylic acid [(1S)-dimethylcarbamoyl-2-phenyl-ethyl]-amide;
- 25 2-chloro-6H-thieno[2,3-b]pyrrole-5-carboxylic acid [(1S)-benzyl-2-(1,1-dioxo-1-thiazolidin-3-yl)-2-oxo-ethyl]-amide;
- 1-((2S)-[(2-chloro-6H-thieno[2,3-b]pyrrole-5-carbonyl)-amino]-3-phenyl-propionyl)-piperidine-4-carboxylic acid ethyl ester;

- 2-bromo-6H-thieno[2,3-b]pyrrole-5-carboxylic acid [(1S)-benzyl-2-(3-hydroxy-azetidin-1-yl)-2-oxo-ethyl]-amide;
- 2-methyl-4H-furo[3,2-b]pyrrole-5-carboxylic acid [(1S)-benzyl-2-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide;
- 5 2-trimethylsilanylethynyl-6H-thieno[2,3-b]pyrrole-5-carboxylic acid [(1S)-benzyl-2-(3-hydroxy-azetidin-1-yl)-2-oxo-ethyl]-amide;
- 2-ethynyl-6H-thieno[2,3-b]pyrrole-5-carboxylic acid [(1S)-benzyl-2-(3-hydroxy-azetidin-1-yl)-2-oxo-ethyl]-amide;
- 2-fluoro-4H-thieno[3,2-b]pyrrole-5-carboxylic acid [(1S)-benzyl-2-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide;
- 10 2-cyano-4H-furo[3,2-b]pyrrole-5-carboxylic acid [(1S)-benzyl-2-(3-hydroxy-azetidin-1-yl)-2-oxo-ethyl]-amide;
- 2-chloro-4H-furo[3,2-b]pyrrole-5-carboxylic acid [(1S)-benzyl-2-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide;
- 15 2-chloro-4H-furo[3,2-b]pyrrole-5-carboxylic acid [(1S)-benzyl-3-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-(2R)-hydroxy-3-oxo-propyl]-amide;
- 1-((2S)-[(2-chloro-6H-thieno[2,3-b]pyrrole-5-carbonyl)-amino]-3-phenyl-propionyl)-piperidine-4-carboxylic acid;
- 3-chloro-4H-thieno[3,2-b]pyrrole-5-carboxylic acid [(1S)-benzyl-2-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide;
- 20 3-chloro-4H-thieno[3,2-b]pyrrole-5-carboxylic acid [(1S)-benzyl-3-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-(2R)-hydroxy-3-oxo-propyl]-amide;
- 3-bromo-4H-thieno[3,2-b]pyrrole-5-carboxylic acid [(1S)-benzyl-2-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide;
- 25 3-bromo-4H-thieno[3,2-b]pyrrole-5-carboxylic acid [(1S)-benzyl-3-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-(2R)-hydroxy-3-oxo-propyl]-amide;
- 2-chloro-4H-thieno[3,2-b]pyrrole-5-carboxylic acid [(1S)-benzyl-3-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-(2R)-hydroxy-3-oxo-propyl]-amide;
- 2-chloro-4H-thieno[3,2-b]pyrrole-5-carboxylic acid [(1S)-benzyl-2-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide;
- 30 3-methyl-4H-thieno[3,2-b]pyrrole-5-carboxylic acid [(1S)-benzyl-2-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide;
- 3-methyl-4H-thieno[3,2-b]pyrrole-5-carboxylic acid [(1S)-benzyl-3-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-(2R)-hydroxy-3-oxo-propyl]-amide;



- 2-cyano-4H-thieno[3,2-b]pyrrole-5-carboxylic acid [(1S)-benzyl-2-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide;
- 2-cyano-4H-furo[3,2-b]pyrrole-5-carboxylic acid [(1S)-benzyl-3-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-(2R)-hydroxy-3-oxo-propyl]-amide;
- 5 3-bromo-4H-furo[3,2-b]pyrrole-5-carboxylic acid [(1S)-benzyl-2-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide;
- 3-bromo-4H-furo[3,2-b]pyrrole-5-carboxylic acid [(1S)-benzyl-3-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-(2R)-hydroxy-3-oxo-propyl]-amide;
- 10 4H-1,7-dithia-4-aza-cyclopenta[a]pentalene-5-carboxylic acid [(1S)-benzyl-3-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-(2R)-hydroxy-3-oxo-propyl]-amide;
- 4H-1,7-dithia-4-aza-cyclopenta[a]pentalene-5-carboxylic acid [(1S)-benzyl-2-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide;
- 2-chloro-3-methyl-4H-thieno[3,2-b]pyrrole-5-carboxylic acid [(1S)-benzyl-2-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide;
- 15 2-chloro-3-methyl-4H-thieno[3,2-b]pyrrole-5-carboxylic acid [(1S)-benzyl-3-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-(2R)-hydroxy-3-oxo-propyl]-amide;
- 2-methylsulfanyl-4H-thieno[3,2-b]pyrrole-5-carboxylic acid [(1S)-benzyl-2-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide;
- 2-Bromo-4H-thieno[3,2-b]pyrrole-5-carboxylic acid [(1S)-benzyl-2-(3-hydroxy-azetidin-1-yl)-2-oxo-ethyl]-amide;
- 20 2-Bromo-4H-thieno[3,2-b]pyrrole-5-carboxylic acid [(1S)-benzyl-2-(1,1-dioxo-1-thiazolidin-3-yl)-2-oxo-ethyl]-amide;
- 2-Bromo-4H-thieno[3,2-b]pyrrole-5-carboxylic acid [(1S)-benzyl-2-morpholin-4-yl-2-oxo-ethyl]-amide;
- 25 2-bromo-4H-thieno[3,2-b]pyrrole-5-carboxylic acid [(1S)-benzyl-2-((3S,4S)-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide;
- 2-bromo-4H-thieno[3,2-b]pyrrole-5-carboxylic acid [(1S)-benzyl-2-((3R,4R)-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide;
- 2-bromo-4H-thieno[3,2-b]pyrrole-5-carboxylic acid [(1S)-benzyl-2-(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-amide; and the pharmaceutically acceptable salts and
- 30 prodrugs thereof, and salts of the prodrugs.

Methods for making the above recited glycogen phosphorylase inhibitors can be found in U.S. provisional patent application number 60/157,148, filed September 30, 1999.

Commonly assigned PCT published applications WO 96/39384 and WO 96/39385 disclose additional glycogen phosphorylase inhibitors that can be used in combination with a neurotensin receptor ligand. Additional preferred glycogen phosphorylase inhibitors include:

- 5            5-chloro-1H-indole-2-carboxylic acid [(1S)-((R)-hydroxy-dimethylcarbamoyl-methyl)-2-phenyl-ethyl]-amide;
- 5,6-dichloro-1H-indole-2-carboxylic acid {(1S)-[(R)-hydroxy-(methoxy-methyl-carbamoyl)-methyl]-2-phenyl-ethyl}-amide;
- 5-chloro-1H-indole-2-carboxylic acid {(1S)-[(R)-hydroxy-(methoxy-methyl-
- 10   carbamoyl)-methyl]-2-phenyl-ethyl}-amide;
- 5-chloro-1H-indole-2-carboxylic acid ((1S)-{(R)-hydroxy-[(2-hydroxy-ethyl)-methyl-carbamoyl]-methyl}-2-phenyl-ethyl)-amide;
- 5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-3-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-(2R)-hydroxy-3-oxo-propyl]-amide;
- 15            5-chloro-1H-indole-2-carboxylic acid {(1S)-[(R)-hydroxy-(methyl-pyridin-2-yl-carbamoyl)-methyl]-2-phenyl-ethyl}-amide;
- 5-chloro-1H-indole-2-carboxylic acid ((1S)-{(R)-hydroxy-[methyl-(2-pyridin-2-yl-ethyl)-carbamoyl]-methyl}-2-phenyl-ethyl)-amide;
- 5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-(2R)-hydroxy-3-(4-methyl-
- 20   piperazin-1-yl)-3-oxo-propyl]-amide hydrochloride;
- 5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-(2R)-hydroxy-3-(3-hydroxy-azetidin-1-yl)-3-oxo-propyl]-amide;
- 5-chloro-1H-indole-2-carboxylic acid ((1S)-benzyl-(2R)-hydroxy-3-isoxazolidin-
- 25   2-yl-3-oxo-propyl)-amide;
- 5-chloro-1H-indole-2-carboxylic acid ((1S)-benzyl-(2R)-hydroxy-3-
- [1,2]oxazinan-2-yl-3-oxo-propyl)-amide;
- 5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-(2R)-hydroxy-3-((3S)-
- hydroxy-pyrrolidin-1-yl)-3-oxo-propyl]-amide;
- 5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-3-((3S,4S)-dihydroxy-
- 30   pyrrolidin-1-yl)-(2R)-hydroxy-3-oxo-propyl]-amide;
- 5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-3-(cis-3,4-dihydroxy-pyrrolidin-1-yl)-(2R)-hydroxy-3-oxo-propyl]-amide;
- 5-chloro-1H-indole-2-carboxylic acid ((1S)-benzyl-(2R)-hydroxy-3-morpholin-
- 4-yl-3-oxo-propyl)-amide;

5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-(3-hydroxyimino-pyrrolidin-1-yl)-2-oxo-ethyl]-amide;

5-chloro-1H-indole-2-carboxylic acid [2-(cis-3,4-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide;

5 5-chloro-1H-indole-2-carboxylic acid [2-((3S,4S)-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide;

5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-(cis-3,4-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide;

10 5-chloro-1H-indole-2-carboxylic acid [2-(1,1-dioxo-thiazolidin-3-yl)-2-oxo-ethyl]-amide;

5-chloro-1H-indole-2-carboxylic acid (2-oxo-2-thiazolidin-3-yl-ethyl)-amide,

5-chloro-1H-indole-2-carboxylic acid [(1S)-(4-fluoro-benzyl)-2-(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-amide;

15 5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-((3RS)-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-amide;

5-chloro-1H-indole-2-carboxylic acid [2-oxo-2-((1RS)-oxo-1-thiazolidin-3-yl)-ethyl]-amide;

5-chloro-1H-indole-2-carboxylic acid [(1S)-(2-fluoro-benzyl)-2-(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-amide;

20 5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-((3S,4S)-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide;

5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-(3-hydroxy-azetidin-1-yl)-2-oxo-ethyl]-amide;

25 5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-(3-hydroxyimino-azetidin-1-yl)-2-oxo-ethyl]-amide;

5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-(4-hydroxyimino-piperidin-1-yl)-2-oxo-ethyl]-amide;

5-chloro-1H-indole-2-carboxylic acid [1-benzyl-2-(3-hydroxypyrrolidin-1-yl)-2-oxo-ethyl]amide;

30 5-chloro-1H-indole-2-carboxylic acid [(1S)-((R)-hydroxy-dimethylcarbamoyl-methyl)-2-phenyl-ethyl]-amide;

5-chloro-1H-indole-2-carboxylic acid [(1S)-((R)-hydroxy-(methoxy-methyl-carbamoyl)-methyl)-2-phenyl-ethyl]-amide;



3-isopropyl-4-(2-chlorophenyl)-1,4-dihydro-1-ethyl-2-methylpyridine; WO 9524391; WO 9709040; WO 9840353; WO 9850359; WO 9731901; EP 884050; and Hoover et al., *J. Med. Chem.*, 1998, 41, 2934-2938.

5 A neurotensin receptor ligand can also be used in combination with an aldose reductase inhibitor. Aldose reductase inhibitors constitute a class of compounds that have become widely known for their utility in treating conditions arising from complications of diabetes, such as diabetic neuropathy and nephropathy. Such compounds are well known to those skilled in the art and are readily identified by standard biological tests. For example, the aldose reductase  
10 inhibitor zopolrestat, 1-phthalazineacetic acid, 3,4-dihydro-4-oxo-3-[[5-(trifluoromethyl)-2-benzothiazolyl]methyl]-, and related compounds are described in U.S. patent 4,939,140.

Aldose reductase inhibitors have been taught for use in lowering lipid levels in mammals. See, for example, U. S. patent 4,492,706 and EP 0 310 931 A2.

15 U. S. patent 5,064,830 discloses the use of certain oxophthalazinyl acetic acid aldose reductase inhibitors, including zopolrestat, for lowering of blood uric acid levels.

Commonly assigned U.S. patent 5,391,551 discloses the use of certain aldose reductase inhibitors, including zopolrestat, for lowering blood lipid levels in  
20 humans. The disclosure teaches that therapeutic utilities derive from the treatment of diseases caused by an increased level of triglycerides in the blood, such diseases include cardiovascular disorders such as thrombosis, arteriosclerosis, myocardial infarction, and angina pectoris. A preferred aldose reductase inhibitor is zopolrestat.

The term aldose reductase inhibitor refers to a compound that inhibits the  
25 bioconversion of glucose to sorbitol, which is catalyzed by the enzyme aldose reductase. Any aldose reductase inhibitor may be used in a combination with a neurotensin receptor ligand. Aldose reductase inhibition is readily determined by those skilled in the art according to standard assays (J. Malone, *Diabetes*, 29:861-864 (1980) "Red Cell Sorbitol, an Indicator of Diabetic Control"). A variety of aldose  
30 reductase inhibitors are described herein; however, other aldose reductase inhibitors useful in certain of the compositions and methods of this invention will be known to those skilled in the art.

The activity of an aldose reductase inhibitor in a tissue can be determined by testing the amount of aldose reductase inhibitor that is required to lower tissue

2025 RELEASE UNDER E.O. 14176

sorbitol (i.e., by inhibiting the further production of sorbitol consequent to blocking aldose reductase) or lower tissue fructose (by inhibiting the production of sorbitol consequent to blocking aldose reductase and consequently the production of fructose.

5           Accordingly, examples of aldose reductase inhibitors useful in certain of the compositions, combinations and methods of the present invention include:

1. 3-(4-bromo-2-fluorobenzyl)-3,4-dihydro-4-oxo-1-phthalazineacetic acid (ponalrestat, US 4,251,528);
2. N[[[(5-trifluoromethyl)-6-methoxy-1-naphthalenyl]thioxomethyl]-N-  
10 methylglycine (tolrestat, US 4,600,724);
3. 5-[(Z,E)- $\beta$ -methylcinnamylidene]-4-oxo-2-thioxo-3-thiazolideneacetic acid (epalrestat, US 4,464,382, US 4,791,126, US 4,831,045);
4. 3-(4-bromo-2-fluorobenzyl)-7-chloro-3,4-dihydro-2,4-dioxo-1(2H)-  
quinazolineacetic acid (zenarestat, US 4,734,419, and 4,883,800);
- 15 5. 2R,4R-6,7-dichloro-4-hydroxy-2-methylchroman-4-acetic acid (US 4,883,410);
6. 2R,4R-6,7-dichloro-6-fluoro-4-hydroxy-2-methylchroman-4-acetic acid (US 4,883,410);
7. 3,4-dihydro-2,8-diisopropyl-3-oxo-2H-1,4-benzoxazine-4-acetic acid (US  
20 4,771,050);
8. 3,4-dihydro-3-oxo-4-[(4,5,7-trifluoro-2-benzothiazolyl)methyl]-2H-1,4-benzothiazine-2-acetic acid (SPR-210, U.S. 5,252,572);
9. N-[3,5-dimethyl-4-[(nitromethyl)sulfonyl]phenyl]-2-methylbenzeneacetamide (ZD5522, U.S. 5,270,342 and U.S. 5,430,060);
- 25 10. (S)-6-fluorospiro[chroman-4,4'-imidazolidine]-2,5'-dione (sorbinil, US 4,130,714);
11. d-2-methyl-6-fluoro-spiro(chroman-4',4'-imidazolidine)-2',5'-dione (US 4,540,704);
12. 2-fluoro-spiro(9H-fluorene-9,4'-imidazolidine)2',5'-dione (US 4,438,272);
- 30 13. 2,7-di-fluoro-spiro(9H-fluorene-9,4'-imidazolidine)2',5'-dione (US 4,436,745, US 4,438,272);
14. 2,7-di-fluoro-5-methoxy-spiro(9H-fluorene-9,4' -imidazolidine)2',5'-dione (US 4,436,745, US 4,438,272);

15. 7-fluoro-spiro(5H-indenol[1,2-b]pyridine-5,3'-pyrrolidine)2,5'-dione (US 4,436,745, US 4,438,272);
16. d-cis-6'-chloro-2',3'-dihydro-2'-methyl-spiro-(imidazolidine-4,4'-4'-H-pyrano(2,3-b)pyridine)-2,5-dione (US 4,980,357);
- 5      17. spiro[imidazolidine-4,5'(6H)-quinoline]2,5-dione-3'-chloro-7,'8'-dihydro-7'-methyl-(5'-cis)(US 5,066,659);
18. (2S,4S)-6-fluoro-2',5'-dioxospiro(chroman-4,4'-imidazolidine)-2-carboxamide (US 5,447,946);
19. 2-[(4-bromo-2-fluorophenyl)methyl]-6-fluorospiro[isoquinoline-4(1H),3'-pyrrolidine]-1,2',3,5'(2H)-tetrone (ARI-509, US 5,037,831);
- 10      20. 3,4-dihydro-3-(5-fluorobenzothiazol-2-ylmethyl)-4-oxophthalazin-1-ylacetic acid;
21. 3-(5,7-difluorobenzothiazol-2-ylmethyl)-3,4-dihydro-4-oxophthalazin-1-ylacetic acid;
- 15      22. 3-(5-chlorobenzothiazol-2-ylmethyl)-3,4-dihydro-4-oxophthalazin-1-ylacetic acid;
23. 3-(5,7-dichlorobenzothiazol-2-ylmethyl)-3,4-dihydro-4-oxophthalazin-1-ylacetic acid;
24. 3,4-dihydro-4-oxo-3-(5-trifluoromethylbenzoxazol-2-ylmethyl)phthalazin-1-ylacetic acid;
- 20      25. 3,4-dihydro-3-(5-fluorobenzoxazol-2-ylmethyl)-4-oxophthalazin-1-ylacetic acid;
26. 3-(5,7-difluorobenzoxazol-2-ylmethyl)-3,4-dihydro-4-oxophthalazin-1-ylacetic acid;
- 25      27. 3-(5-chlorobenzoxazol-2-ylmethyl)-3,4-dihydro-4-oxophthalazin-1-ylacetic acid;
28. 3-(5,7-dichlorobenzoxazol-2-ylmethyl)-3,4-dihydro-4-oxophthalazin-1-ylacetic acid;
29. zopolrestat; 1-phthalazineacetic acid, 3,4-dihydro-4-oxo-3-[[5-(trifluoromethyl)-2-benzothiazolyl]methyl]-; and the pharmaceutically acceptable salts and prodrugs thereof, and salts of the prodrugs.
- 30

Procedures for making the aldose reductase inhibitors 20-29 are disclosed in PCT publication number WO 99/26659.

A neurotensin receptor ligand can also be used in combination with a sorbitol dehydrogenase inhibitor. Sorbitol dehydrogenase inhibitors lower fructose levels and have been used to treat or prevent diabetic complications such as neuropathy, retinopathy, nephropathy, cardiomyopathy, microangiopathy, and macroangiopathy. U.S. patent numbers 5,728,704 and 5,866,578 disclose compounds and methods for treating diabetic complications by inhibiting the enzyme sorbitol dehydrogenase. The compounds disclosed in these patents and other sorbitol dehydrogenase inhibitors can be used in the present invention in combination with a neurotensin receptor ligand.

10 A neurotensin receptor ligand can also be used in combination with a glucocorticoid receptor antagonist. The glucocorticoid receptor (GR) is present in glucocorticoid responsive cells where it resides in the cytosol in an inactive state until it is stimulated by an agonist. Upon stimulation the glucocorticoid receptor translocates to the cell nucleus where it specifically interacts with DNA and/or  
15 protein(s) and regulates transcription in a glucocorticoid responsive manner. Two examples of proteins that interact with the glucocorticoid receptor are the transcription factors, API and NF $\kappa$ - $\beta$ . Such interactions result in inhibition of API- and NF $\kappa$ - $\beta$ -mediated transcription and are believed to be responsible for the anti-inflammatory activity of endogenously administered glucocorticoids. In addition,  
20 glucocorticoids may also exert physiologic effects independent of nuclear transcription. Biologically relevant glucocorticoid receptor agonists include cortisol and corticosterone. Many synthetic glucocorticoid receptor agonists exist including dexamethasone, prednisone and prednisilone. By definition, glucocorticoid receptor antagonists bind to the receptor and prevent glucocorticoid receptor  
25 agonists from binding and eliciting GR mediated events, including transcription. RU486 is an example of a non-selective glucocorticoid receptor antagonist. GR antagonists can be used in the treatment of diseases associated with an excess or a deficiency of glucocorticoids in the body. As such, they may be used to treat the following: obesity, diabetes, cardiovascular disease, hypertension, Syndrome X,  
30 depression, anxiety, glaucoma, human immunodeficiency virus (HIV) or acquired immunodeficiency syndrome (AIDS), neurodegeneration (for example, Alzheimer's and Parkinson's), cognition enhancement, Cushing's Syndrome, Addison's Disease, osteoporosis, frailty, inflammatory diseases (such as osteoarthritis, rheumatoid arthritis, asthma and rhinitis), adrenal dysfunction, viral infection,



5            Examples of preferred glucocorticoid receptor antagonists that can be used in combination with a neurotensin receptor ligand can be found in U.S. provisional patent application number 60/132,130.

In addition, a neurotensin receptor ligand can be used in combination with a thyromimetic. It is generally accepted that thyroid hormones, specifically, biologically active iodothyronines, are critical to normal development and to maintaining metabolic homeostasis. Thyroid hormones stimulate the metabolism of cholesterol to bile acids and enhance the lipolytic responses of fat cells to other hormones. U.S. patent numbers 4,766,121; 4,826,876; 4,910,305; and 5,061,798 disclose certain thyroid hormone mimetics (thyromimetics), namely, 3,5-dibromo-3'-[6-oxo-3(1H)-pyridazinylmethyl]-thyronines. U.S. patent number 5,284,971 discloses certain thyromimetic cholesterol lowering agents, namely, 4-(3-cyclohexyl-4-hydroxy or -methoxy phenylsulfonyl)-3,5 dibromo-phenylacetic compounds. U.S. patent numbers 5,401,772; 5,654,468; and 5,569,674 disclose certain thyromimetics that are lipid lowering agents, namely, heteroacetic acid derivatives. In addition, certain oxamic acid derivatives of thyroid hormones are known in the art. For example, N. Yokoyama, *et al.* in an article published in the *Journal of Medicinal Chemistry*, **38 (4)**: 695-707 (1995) describe replacing a -CH<sub>2</sub> group in a naturally occurring metabolite of T<sub>3</sub> with an -NH group resulting in -HNCOCO<sub>2</sub>H. Likewise, R.E. Steele *et al.* in an article published in International Congressional Service (*Atherosclerosis X*) **1066**: 321-324 (1995) and Z.F. Stephan *et al.* in an article published in *Atherosclerosis*, **126**: 53-63 (1996), describe certain oxamic acid derivatives that are useful as lipid-lowering thyromimetic agents and are devoid of undesirable cardiac activities.

In addition, a neurotensin receptor ligand can be administered in combination with other pharmaceutical agents such as cholesterol biosynthesis inhibitors and cholesterol absorption inhibitors, especially HMG-CoA reductase inhibitors and HMG-CoA synthase inhibitors, HMG-CoA reductase and synthase gene expression inhibitors, CETP inhibitors, bile acid sequesterants, fibrates, ACAT inhibitors, squalene synthetase inhibitors, anti-oxidants and niacin. A neurotensin receptor ligand may also be administered in combination with naturally occurring compounds that act to lower plasma cholesterol levels. These naturally occurring compounds are commonly called nutraceuticals and include, for example, garlic extract, Benecol®, and niacin.

Specific cholesterol absorption inhibitors and cholesterol biosynthesis inhibitors are described in detail below. Additional cholesterol absorption inhibitors are known to those skilled in the art and are described, for example, in WO 94/00480.

Any HMG-CoA reductase inhibitor may be employed as an additional compound in the combination therapy aspect of the present invention. The term HMG-CoA reductase inhibitor refers to a compound that inhibits the biotransformation of hydroxymethylglutaryl-coenzyme A to mevalonic acid as catalyzed by the enzyme HMG-CoA reductase. Such inhibition may be determined readily by one of skill in the art according to standard assays (e.g., *Methods of Enzymology*, 71: 455-509 (1981); and the references cited therein). A variety of these compounds are described and referenced below. U.S. patent number 4,231,938 discloses certain compounds isolated after cultivation of a microorganism belonging to the genus *Aspergillus*, such as lovastatin. Also, U.S. patent number 4,444,784 discloses synthetic derivatives of the aforementioned compounds, such as simvastatin. Additionally, U.S. patent number 4,739,073 discloses certain substituted indoles, such as fluvastatin. Further, U.S. patent number 4,346,227 discloses ML-236B derivatives, such as pravastatin. In addition, EP 491,226 teaches certain pyridyldihydroxyheptenoic acids, such as rivastatin. Also, U.S. patent number 4,647,576 discloses certain 6-[2-(substituted-pyrrol-1-yl)-alkyl]-pyran-2-ones such as atorvastatin. Other HMG-CoA reductase inhibitors will be known to those skilled in the art. Examples of marketed products containing HMG-CoA reductase inhibitors that can be used in combination with compounds of the present invention include Baycol®, Lescol®, Lipitor®, Mevacor®, Pravachol® and Zocor®.

Any HMG-CoA synthase inhibitor may be used as the second compound in the combination therapy aspect of this invention. The term HMG-CoA synthase inhibitor refers to a compound which inhibits the biosynthesis of hydroxymethylglutaryl-coenzyme A from acetyl-coenzyme A and acetoacetyl-coenzyme A, catalyzed by the enzyme HMG-CoA synthase. Such inhibition may be determined readily by one of skill in the art according to standard assays (e.g., *Methods of Enzymology*, **35**: 155-160 (1975); and *Methods of Enzymology*, **110**: 19-26 (1985); and the references cited therein). A variety of these compounds are described and referenced below. U.S. patent number 5,120,729 discloses certain beta-lactam derivatives. U.S. patent number 5,064,856 discloses certain spiro-lactone derivatives prepared by culturing the microorganism MF5253. U.S. patent number 4,847,271 discloses certain oxetane compounds such as 11-(3-hydroxymethyl-4-oxo-2-oxetanyl)-3,5,7-trimethyl-2,4-undecadienoic acid derivatives. Other HMG-CoA synthase inhibitors will be known to those skilled in the art.

Any compound that decreases HMG-CoA reductase gene expression may be used as an additional compound in the combination therapy aspect of this invention. These agents may be HMG-CoA reductase transcription inhibitors that block the transcription of DNA or translation inhibitors that prevent translation of mRNA coding for HMG-CoA reductase into protein. Such inhibitors may either affect transcription or translation directly, or may be biotransformed into compounds that have the aforementioned attributes by one or more enzymes in the cholesterol biosynthetic cascade or may lead to the accumulation of an isoprene metabolite that has the aforementioned activities. Such regulation is readily determined by those skilled in the art according to standard assays (*Methods of Enzymology*, **110**: 9-19 1985).

Several such compounds are described and referenced below however other inhibitors of HMG-CoA reductase gene expression will be known to those skilled in the art. U.S. Patent Number 5,041,432 discloses certain 15-substituted lanosterol derivatives. Other oxygenated sterols that suppress the biosynthesis of HMG-CoA reductase are discussed by E.I. Mercer (*Prog. Lip. Res.*, **32**:357-416 1993).

Any compound having activity as a CETP inhibitor can serve as the second compound in the combination therapy aspect of the instant invention. The term CETP inhibitor refers to compounds that inhibit the cholesteryl ester transfer protein (CETP) mediated transport of various cholesteryl esters and triglycerides from HDL to LDL and VLDL. A variety of these compounds are

described and referenced below however other CETP inhibitors will be known to those skilled in the art. U.S. patent number 5,512,548 discloses certain polypeptide derivatives having activity as CETP inhibitors, while certain CETP-inhibitory rosenonolactone derivatives and phosphate-containing analogs of cholesteryl ester are disclosed in *J. Antibiot.*, **49**(8): 815-816 (1996), and *Bioorg. Med. Chem. Lett.*; **6**:1951-1954 (1996), respectively. Other CETP inhibitors that can be used in combination with a neurotensin receptor ligand are disclosed in WO 99/20302, EP 796846, EP818197, EP 818448, WO 99/14204, WO 99/41237, WO 95/04755, WO 96/15141, WO 96/05227, DE 19704244, DE19741051, DE 19741399, DE 19704243, DE 19709125, DE 19627430, DE 19832159, DE 19741400, JP 11049743, and JP 09059155. Preferred CETP inhibitors that can be used in combination with a neurotensin receptor ligand include:

[2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;

[2S,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-methoxymethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;

[2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid 2-hydroxy-ethyl ester;

[2S,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-cyclopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;

[2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;

[2S,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-cyclopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid propyl ester;

[2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid propyl ester;

[2S,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-isopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;

- [2S,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-6-chloro-2-cyclopropyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;
- [2S,4S] 2-cyclopropyl-4-[(3,5-dichloro-benzyl)-methoxycarbonyl-amino]-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;
- 5 [2S,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-cyclopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid tert-butyl ester;
- [2S,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-cyclopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid
- 10 isopropyl ester;
- [2S,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-cyclobutyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;
- [2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;
- 15 [2S,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-methoxymethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;
- [2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid 2-hydroxy-ethyl
- 20 ester;
- [2S,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-cyclopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;
- [2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;
- 25 [2S,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-cyclopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid propyl ester;
- [2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid propyl ester; and
- 30 the pharmaceutically acceptable salts and prodrugs thereof, and the salts of the prodrugs.

Any compound having activity as a squalene synthetase inhibitor can serve as an additional compound in the combination therapy aspect of the instant invention. The term squalene synthetase inhibitor refers to a compound that inhibits the condensation of two molecules of farnesylpyrophosphate to form squalene, a reaction that is catalyzed by the enzyme squalene synthetase. Such inhibition is readily determined by those skilled in the art according to standard methodology (*Methods of Enzymology*, **15**:393-454 (1969); and *Methods of Enzymology*, **110**: 359-373 (1985); and references cited therein). A summary of squalene synthetase inhibitors has been compiled in *Curr. Op. Ther. Patents*, 861-4, (1993). European patent application publication number 0 567 026 A1 discloses certain 4,1-benzoxazepine derivatives as squalene synthetase inhibitors and their use in the treatment of hypercholesterolemia and as fungicides. European patent application publication number 0 645 378 A1 discloses certain seven- or eight-membered heterocycles as squalene synthetase inhibitors and their use in the treatment and prevention of hypercholesterolemia and fungal infections. European patent application publication number 0 645 377 A1 discloses certain benzoxazepine derivatives as squalene synthetase inhibitors useful for the treatment of hypercholesterolemia or coronary sclerosis. European patent application publication number 0 611 749 A1 discloses certain substituted amic acid derivatives useful for the treatment of arteriosclerosis. European patent application publication number 0 705 607 A2 discloses certain condensed seven- or eight-membered heterocyclic compounds useful as antihypertriglyceridemic agents. PCT publication WO 96/09827 discloses certain combinations of cholesterol absorption inhibitors and cholesterol biosynthesis inhibitors including benzoxazepine derivatives and

Any compound having activity as a squalene synthetase inhibitor can serve as an additional compound in the combination therapy aspect of the instant invention. The term squalene synthetase inhibitor refers to a compound that inhibits the condensation of two molecules of farnesylpyrophosphate to form squalene, a reaction that is catalyzed by the enzyme squalene synthetase. Such inhibition is readily determined by those skilled in the art according to standard methodology (*Methods of Enzymology*, **15**:393-454 (1969); and *Methods of Enzymology*, **110**: 359-373 (1985); and references cited therein). A summary of squalene synthetase inhibitors has been compiled in *Curr. Op. Ther. Patents*, 861-4, (1993). European patent application publication number 0 567 026 A1 discloses certain 4,1-benzoxazepine derivatives as squalene synthetase inhibitors and their use in the treatment of hypercholesterolemia and as fungicides. European patent application publication number 0 645 378 A1 discloses certain seven- or eight-membered heterocycles as squalene synthetase inhibitors and their use in the treatment and prevention of hypercholesterolemia and fungal infections. European patent application publication number 0 645 377 A1 discloses certain benzoxazepine derivatives as squalene synthetase inhibitors useful for the treatment of hypercholesterolemia or coronary sclerosis. European patent application publication number 0 611 749 A1 discloses certain substituted amic acid derivatives useful for the treatment of arteriosclerosis. European patent application publication number 0 705 607 A2 discloses certain condensed seven- or eight-membered heterocyclic compounds useful as antihypertriglyceridemic agents. PCT publication WO 96/09827 discloses certain combinations of cholesterol absorption inhibitors and cholesterol biosynthesis inhibitors including benzoxazepine derivatives and

5

10

15

20

30

usually secreted in vast excess of the amounts necessary for the breakdown of fats in the upper small intestine. Because pancreatic lipase is the primary enzyme required for the absorption of dietary triglycerides, inhibitors have utility in the treatment of obesity and the other related conditions.

5           Gastric lipase is an immunologically distinct lipase that is responsible for approximately 10 to 40% of the digestion of dietary fats. Gastric lipase is secreted in response to mechanical stimulation, ingestion of food, the presence of a fatty meal or by sympathetic agents. Gastric lipolysis of ingested fats is of physiological importance in the provision of fatty acids needed to trigger pancreatic lipase activity in the  
10 intestine and is also of importance for fat absorption in a variety of physiological and pathological conditions associated with pancreatic insufficiency. See, for example, C.K. Abrams, et al., *Gastroenterology*, **92**, 125 (1987).

A variety of lipase inhibitors are known to one of ordinary skill in the art. However, in the practice of certain of the methods, pharmaceutical compositions and  
15 kits of the instant invention, generally preferred lipase inhibitors are those inhibitors that are selected from the group consisting of lipstatin, tetrahydrolipstatin (orlistat), FL-386, WAY-121898, Bay-N-3176, valilactone, esterastin, ebelactone A, ebelactone B and RHC 80267.

The pancreatic lipase inhibitors lipstatin, 2S, 3S, 5S, 7Z, 10Z)-5-[(S)-2-  
20 formamido-4-methyl-valeryloxy]-2-hexyl-3-hydroxy-7,10-hexadecanoic acid lactone, and tetrahydrolipstatin (orlistat), 2S, 3S, 5S)-5-[(S)-2-formamido-4-methyl-valeryloxy]-2-hexyl-3-hydroxy-hexadecanoic acid lactone, and the variously substituted N-formylleucine derivatives and stereoisomers thereof, are disclosed in U.S. patent number 4,598,089.

25           The pancreatic lipase inhibitor FL-386, 1-[4-(2-methylpropyl)cyclohexyl]-2-[(phenylsulfonyl)oxy]-ethanone, and the variously substituted sulfonate derivatives related thereto, are disclosed in U.S. patent number 4,452,813.

The pancreatic lipase inhibitor WAY-121898, 4-phenoxyphenyl-4-methylpiperidin-1-yl-carboxylate, and the various carbamate esters and  
30 pharmaceutically acceptable salts related thereto, are disclosed in U.S. patent numbers 5,512,565; 5,391,571 and 5,602,151.

The lipase inhibitor Bay-N-3176, N-3-trifluoromethylphenyl-N'-3-chloro-4'-trifluoromethylphenylurea, and the various urea derivatives related thereto, are disclosed in U.S. patent number 4,405,644.



The lipase inhibitor esteracin, and certain processes for the preparation thereof by the microbial cultivation of *Streptomyces* strain ATCC 31336, are disclosed in U.S. patent numbers 4,189,438 and 4,242,453.

The pancreatic lipase inhibitors ebelactone A and ebelactone B, and a process for the preparation thereof by the microbial cultivation of *Actinomycetes* strain MG7-G1, are disclosed in Umezawa, et al., *J. Antibiotics*, **33**, 1594-1596 (1980). The use of ebelactones A and B in the suppression of monoglyceride formation is disclosed in Japanese Kokai 08-143457, published June 4, 1996.

A glucosidase inhibitor inhibits the enzymatic hydrolysis of complex carbohydrates by glycoside hydrolases, for example amylase or maltase, into bioavailable simple sugars, for example, glucose. The rapid metabolic action of glucosidases, particularly following the intake of high levels of carbohydrates, results in a state of alimentary hyperglycemia which, in adipose or diabetic subjects, leads to enhanced secretion of insulin, increased fat synthesis and a reduction in fat degradation. Following such hyperglycemias, hypoglycemia frequently occurs, due to the augmented levels of insulin present. Additionally, it is known that both hypoglycemias and chyme remaining in the stomach promotes the production of gastric juice, which initiates or favors the development of gastritis or duodenal ulcers. Accordingly, glucosidase inhibitors are known to have utility in accelerating the passage of carbohydrates through the stomach and inhibiting the absorption of glucose from the intestine. Furthermore, the conversion of carbohydrates into lipids of fatty tissue and the subsequent incorporation of alimentary fat into fatty tissue deposits is accordingly reduced or delayed, with the concomitant benefit of reducing or preventing the deleterious abnormalities resulting therefrom.

In combination with a neurotensin receptor ligand, any glucosidase inhibitor may be employed; however, a generally preferred glucosidase inhibitor comprises an amylase inhibitor. An amylase inhibitor is a glucosidase inhibitor that inhibits the enzymatic degradation of starch or glycogen into maltose. The inhibition of such enzymatic degradation is beneficial in reducing amounts of bioavailable sugars, including glucose and maltose, and the concomitant deleterious conditions resulting therefrom.

A variety of glucosidase and amylase inhibitors are known to one of ordinary skill in the art. However, in the practice of the methods and pharmaceutical compositions of the instant invention, generally preferred glucosidase inhibitors are those inhibitors that are selected from the group consisting of acarbose, adiposine, voglibose, miglitol, emiglitate, MDL-25637, camiglibose, tendamistate, AI-3688, trestatin, pradimicin-Q and salbostatin.

The glucosidase inhibitor acarbose, O-4,6-dideoxy-4-[[[(1S,4R,5S,6S)-4,5,6-trihydroxy-3-(hydroxymethyl)-2-cyclohexen-1-yl]amino]- $\alpha$ -glucopyranosyl-(1 $\rightarrow$ 4)-O- $\alpha$ -D-glucopyranosyl-(1 $\rightarrow$ 4)-D-glucose, the various amino sugar derivatives related thereto and a process for the preparation thereof by the microbial cultivation of *Actinoplanes* strains SE 50 (CBS 961.70), SB 18 (CBS 957.70), SE 82 (CBS 615.71), SE 50/13 (614.71) and SE 50/110 (674.73) are disclosed in U.S. patent numbers 4,062,950 and 4,174,439 respectively.

The glucosidase inhibitor adiposine, consisting of adiposine forms 1 and 2, is disclosed in U.S. patent number 4,254,256. Additionally, a process for the preparation and purification of adiposine is disclosed in Namiki et al., *J. Antibiotics*, **35**, 1234-1236 (1982).

The glucosidase inhibitor voglibose, 3,4-dideoxy-4-[[2-hydroxy-1-(hydroxymethyl)ethyl]amino]-2-C-(hydroxymethyl)-D-epi-inositol, and the various N-substituted pseudo-aminosugars related thereto, are disclosed in U.S. patent number 4,701,559.

The glucosidase inhibitor miglitol, (2R,3R,4R,5S)-1-(2-hydroxyethyl)-2-(hydroxymethyl)-3,4,5-piperidinetriol, and the various 3,4,5-trihydroxypiperidines related thereto, are disclosed in U.S. patent number 4,639,436.

The glucosidase inhibitor emiglitate, ethyl p-[2-[(2R,3R,4R,5S)-3,4,5-trihydroxy-2-(hydroxymethyl)piperidino]ethoxy]-benzoate, the various derivatives

related thereto and pharmaceutically acceptable acid addition salts thereof, are disclosed in U.S. patent number 5,192,772.

The glucosidase inhibitor MDL-25637, 2,6-dideoxy-7-O- $\beta$ -D-glucopyrano-syl-2,6-imino-D-glycero-L-gluco-heptitol, the various homodisaccharides related thereto  
5 and the pharmaceutically acceptable acid addition salts thereof, are disclosed in U.S. patent number 4,634,765.

The glucosidase inhibitor camiglibose, methyl 6-deoxy-6-[(2R,3R,4R,5S)-3,4,5-trihydroxy-2-(hydroxymethyl)piperidino]- $\alpha$ -D-glucopyranoside sesquihydrate, the deoxy-nojirimycin derivatives related thereto, the various pharmaceutically acceptable  
10 salts thereof and synthetic methods for the preparation thereof, are disclosed in U.S. patent numbers 5,157,116 and 5,504,078.

The amylase inhibitor tendamistat, the various cyclic peptides related thereto and processes for the preparation thereof by the microbial cultivation of *Streptomyces tendae* strains 4158 or HAG 1226, are disclosed in U.S. Patent Number 4,451,455.

The amylase inhibitor AI-3688, the various cyclic polypeptides related thereto,  
15 and a process for the preparation thereof by the microbial cultivation of *Streptomyces aureofaciens* strain FH 1656, are disclosed in U.S. patent number 4,623,714.

The amylase inhibitor trestatin, consisting of a mixture of trestatin A, trestatin B and trestatin C, the various trehalose-containing aminosugars related thereto and a  
20 process for the preparation thereof by the microbial cultivation of *Streptomyces dimorphogenes* strains NR-320-OM7HB and NR-320-OM7HBS, are disclosed in U.S. patent number 4,273,765.

The glucosidase inhibitor pradimicin-Q and a process for the preparation thereof by the microbial cultivation of *Actinomadura verrucospora* strains R103-3 or  
25 A10102, are disclosed in U.S. patent numbers 5,091,418 and 5,217,877, respectively.

The glycosidase inhibitor salbostatin, the various pseudosaccharides related thereto, the various pharmaceutically acceptable salts thereof and a process for the preparation thereof by the microbial cultivation of *Streptomyces albus* strain ATCC  
30 21838, are disclosed in U.S. patent number 5,091,524.

Preferred lipase inhibitors comprise compounds selected from the group consisting of lipstatin, tetrahydrolipstatin, FL-386, WAY-121898, Bay-n-3176, valilactone, esteracin, ebelactone A, ebelactone B, RHC 80267, stereoisomers

thereof, and pharmaceutically acceptable salts of said compounds and stereoisomers. The compound tetrahydrolipstatin is especially preferred.

Preferred glucosidase inhibitors comprise compounds selected from the group consisting of acarbose, adiposine, voglibose, miglitol, emiglitate, MDL-25637, 5 camiglibose, pradimicin-Q, and salbostatin. An especially preferred glucosidase inhibitor is acarbose. Especially preferred glucosidase inhibitors further comprise amylase inhibitors that are selected from the group consisting of tendamistate, Al-3688 and trestatin.

In addition, the present invention includes the use of a neurotensin receptor 10 ligand in combination with apo B secretion/MTP inhibitors.

A variety of apo B secretion/MTP inhibitors are known to one of ordinary skill in the art. Although any apo B secretion/MTP inhibitor may be used in the practice of the methods and pharmaceutical compositions of the instant invention, generally preferred apo B secretion/MTP inhibitors include those compounds that are disclosed 15 in, for example, European patent application publication numbers EP 643057, EP 719763, EP 753517, EP 764647, EP 765878, EP 779276, EP 779279, EP 799828, EP 799829, EP 802186, EP 802188, EP 802192, and EP 802197; PCT Application Publication Numbers WO 96/13499, WO 96/33193, WO 96/40640, WO 97/26240, WO 97/43255, WO 97/43257, WO 98/16526 and WO 98/23593; and U.S. patent 20 numbers 5,595,872; 5,646,162; 5,684,014; 5,712,279; 5,739,135 and 5,789,197.

Especially preferred apo-B secretion/MTP inhibitors are those biphenyl-2-carboxylic acid-tetrahydroisoquinolin-6-yl amide derivatives disclosed in PCT application publication numbers WO 96/40640 and WO 98/23593. Especially preferred apo B secretion/MTP inhibitors disclosed in PCT application publication 25 numbers WO 96/40640 and WO 98/23593, and useful in the methods and pharmaceutical compositions of the present invention, are 4'-trifluoromethyl-biphenyl-2-carboxylic acid-[2-(1H-[1,2,4]triazol-3-ylmethyl)-1,2,3,4-tetrahydroisoquin-6-yl]-amide and 4'-trifluoromethyl-biphenyl-2-carboxylic acid-[2-(acetylaminoethyl)-1,2,3,4-tetrahydroisoquinolin-6-yl]-amide.

Another especially preferred class of apo B secretion/MTP inhibitors is 30 disclosed in U.S. patent numbers 5,595,872; 5,721,279; 5,739,135 and 5,789,197.

Especially preferred apo B secretion/MTP inhibitors disclosed in U.S. patent numbers 5,595,872; 5,721,279; 5,739,135 and 5,789,197 and useful in the methods and pharmaceutical compositions of the present invention, are 9-(4-{4-

[4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-piperidin-1-yl)-butyl-9H-fluorene-9-carboxylic acid-(2,2,2-trifluoroethyl)-amide and 9-{4-[4-(2-benzothiazol-2-yl-benzoylamino)-piperidin-1-yl]-butyl}-9H-fluorene-9-carboxylic acid-(2,2,2-trifluoroethyl)-amide.

5 Another class of especially preferred apo B secretion/MTP inhibitors is disclosed in PCT application publication number WO 98/16526.

Especially preferred apo B secretion/MTP inhibitors disclosed in PCT application publication number WO 98/16526, and useful in the methods and pharmaceutical compositions of the present invention, are [11a-R]-8-[(4-cyanophenyl)methoxy]-2-cyclopentyl-7-(prop-2-enyl)-2,3,11,11a-tetrahydro-6H-pyrazino[1,2b]isoquinoline-1,4-dione and [11a-R]-cyclopentyl-7-(prop-2-enyl)-8-  
10 [(pyridin-2-yl)methoxy]-2,3,11,11a-tetrahydro-6H-pyrazino[1,2b]isoquinoline-1,4-dione.

Another especially preferred class of apo B secretion/MTP inhibitors is  
15 disclosed in U.S. patent number 5,684,014.

An especially preferred apo B secretion/MTP inhibitor disclosed in U.S. patent number 5,684,014, and useful in certain of the methods and pharmaceutical compositions of the present invention, is 2-cyclopentyl-2-[4-(2,4-dimethyl-pyrido[2,3-b]indol-9-ylmethyl)-phenyl]-N-(2-hydroxy-1-phenyl-ethyl)-acetamide.

20 Yet another class of especially preferred apo B secretion/MTP inhibitors is disclosed in U.S. patent number 5,646,162.

An especially preferred apo B secretion/MTP inhibitor disclosed in U.S. patent number 5,646,162 and useful in the methods and pharmaceutical compositions of the present invention, is 2-cyclopentyl-N-(2-hydroxy-1-phenylethyl)-2-[4-(quinolin-2-ylmethoxy)-phenyl]-acetamide.  
25

Additional apo B secretion/MTP inhibitors that can be used in combination with a neurotensin receptor ligand are disclosed in U.S. provisional patent application number 60/164,803. Examples of specific preferred apo B secretion/MTP inhibitors disclosed in that application include:

30 7-amino-quinoline-3-carboxylic acid ethyl ester;  
7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid ethyl ester;  
7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid;

- 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid  
(dipyridin-2-yl-methyl)-amide;
- 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid  
(dipyridin-2-yl-methyl)-amide, ethanesulfonate;
- 5 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid  
(dipyridin-2-yl-methyl)-amide, bis-ethanesulfonate;
- 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid (phenyl-  
pyridin-2-yl-methyl)-amide;
- 10 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid (phenyl-  
pyridin-2-yl-methyl)-amide, ethanesulfonate;
- (S)-7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid  
(phenyl-pyridin-2-yl-methyl)-amide;
- (S)-7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid  
(phenyl-pyridin-2-yl-methyl)-amide, ethanesulfonate;
- 15 (S)-7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid  
(phenyl-pyridin-2-yl-methyl)-amide, bis-ethanesulfonate;
- (R)-7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid  
(phenyl-pyridin-2-yl-methyl)-amide;
- (R)-7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid  
20 (phenyl-pyridin-2-yl-methyl)-amide, ethanesulfonate;
- 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid (phenyl-  
pyridin-2-yl-methyl)-amide, bis-ethanesulfonate;
- 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid (1-  
carbamoyl-2-phenyl-ethyl)-amide;
- 25 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid  
(carbamoyl-phenyl-methyl)-amide;
- 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid  
propylamide;
- 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid (2,2,2-  
30 trifluoro-ethyl)-amide;
- 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid (1-  
methyl-1-phenyl-ethyl)-amide;
- 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid  
cyclopentylamide;

- 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid (1-phenyl-propyl)-amide;  
 (R)-7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid (1-phenyl-ethyl)-amide, ethanesulfonate;
- 5 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid (1-phenyl-ethyl)-amide;  
 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid (1-pyridin-2-yl-propyl)-amide;  
 (R)-7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid (1-10 pyridin-2-yl-propyl)-amide;  
 (R)-7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid (1-pyridin-2-yl-propyl)-amide, ethanesulfonate;  
 (S)-7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid (1-pyridin-2-yl-propyl)-amide;
- 15 (S)-7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid (1-pyridin-2-yl-propyl)-amide ethanesulfonate;  
 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid (1-pyridin-2-yl-propyl)-amide, ethanesulfonate;  
 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid (pyridin-20 2-ylmethyl)-amide, ethanesulfonate;  
 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid (2-pyridin-2-yl-ethyl)-amide, ethanesulfonate;  
 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid ethylamide, ethanesulfonate;
- 25 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid butylamide, ethanesulfonate;  
 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid (thiophen-2-ylmethyl)-amide, ethanesulfonate;  
 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid (1-30 methyl-1-pyridin-2-yl-ethyl)-amide;  
 (S)-7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid (1-pyridin-2-yl-ethyl)-amide;  
 (R)-7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid (1-pyridin-2-yl-ethyl)-amide ethanesulfonate;

20044274 04440 9271360

- 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid (1-pyridin-2-yl-ethyl)-amide;  
 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid (1-pyridin-2-yl-ethyl)-amide ethanesulfonate;  
 5 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid amide;  
 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid benzylamide;  
 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid 4-methoxy-benzylamide;  
 10 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid 4-chloro-benzylamide;  
 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid 4-methyl-benzylamide;  
 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid  
 15 cyclopropylmethyl-amide;  
 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid 4-fluoro-benzylamide;  
 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid  
 isopropyl-amide;  
 20 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid benzhydryl-amide;  
 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid cyclopropylamide;  
 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid [1-(4-  
 25 fluoro-phenyl)-ethyl]-amide;  
 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid 3-methyl-benzylamide;  
 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid 3-methoxy-benzylamide;  
 30 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid 3-chloro-benzylamide;  
 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid 2-fluoro-benzylamide;

160000-324300



7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid 3-fluoro-benzylamide;

7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid 2-methyl-benzylamide;

5 7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid 2-methoxy-benzylamide;

7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid 2-chloro-benzylamide;

10 4'-trifluoromethyl-biphenyl-2-carboxylic acid [3-(pyrrolidine-1-carbonyl)-quinolin-7-yl]-amide;

4'-trifluoromethyl-biphenyl-2-carboxylic acid [3-(morpholine-4-carbonyl)-quinolin-7-yl]-amide;

7-[(4'-trifluoromethyl-biphenyl-2-carbonyl)-amino]-quinoline-3-carboxylic acid diethylamide;

15 4'-trifluoromethyl-biphenyl-2-carboxylic acid [3-(piperidine-1-carbonyl)-quinolin-7-yl]-amide; and the pharmaceutically acceptable salts and prodrugs thereof, and salts of the prodrugs.

In addition, a neurotensin receptor ligand can be used in combination with one or more additional compounds that are neurotensin receptor ligands such as those set forth above.

20 Some abbreviations used in this application are defined below:

	GTP	Guanosine 5'-triphosphate
	GDP	Guanosine 5'-diphosphate
	DTT	Dithiothreitol
25	PEI	Polyethylene imine
	FLIPR	Flouresence imaging plate reader
	DMSO	Dimethylsulfoxide
	HEPES	(N-[2-Hydroxyethyl]piperazine-N'-[2-ethanesulfonic acid])
	NT	Neurotensin
30	BSA	Bovine serum albumin
	PBS	Phosphate buffered saline
	EDTA	Ethylenediaminetetraacetic acid
	EGTA	Ethylene glycol-bis( $\beta$ -aminoethyl ether) N,N,N',N'-tetraacetic acid

All documents cited herein are hereby incorporated by reference.

The examples presented below are intended to illustrate particular embodiments of the invention, and are not intended to limit the scope of the specification, including the claims, in any manner.

5

# Examples

## SATURATION BINDING: GTP $\gamma$ [<sup>35</sup>S] BINDING

Agonist-stimulated GTP $\gamma$ [<sup>35</sup>S] binding is a reliable method for the determination of agonist efficacy and potency at G-protein-coupled receptors (GPCRs). GDP binds preferentially over GTP at unactivated heterotrimeric G protein alpha subunits. When an agonist activates a GPCR, the GDP is displaced by GTP (or GTP analogs) causing dissociation of the alpha subunit from the beta and gamma subunits of the heterotrimeric G protein. To measure agonist efficacy and potency GTP $\gamma$ [<sup>35</sup>S] is used instead of GTP.

Well			STOCK	FINAL
	-	+	[GTP $\gamma$ [ <sup>35</sup> S]]	[GTP $\gamma$ [ <sup>35</sup> S]]
A	1,2	3,4	1 nM	0.05 nM
B	1,2	3,4	2 nM	0.1 nM
C	1,2	3,4	5 nM	0.25 nM
D	1,2	3,4	10 nM	0.5 nM
E	1,2	3,4	15 nM	0.75 nM
F	1,2	3,4	20 nM	1 nM
G	1,2	3,4	100 nM	5 nM
H	1,2	3,4	200 nM	10 nM

25

Specific Activity of GTP $\gamma$ [<sup>35</sup>S] is approximately 1250 Ci/mmol. The useful life of GTP $\gamma$ [<sup>35</sup>S] is about one month. Radioactivity calculators that are well known to those skilled in the art can be used to determine the actual concentration of stock GTP $\gamma$ [<sup>35</sup>S].

30

### **To each well add in order:**

20  $\mu$ l binding buffer to "-" wells

20  $\mu$ l of neurotensin in binding buffer (1  $\mu$ M final) to "+" wells

10  $\mu$ l appropriate concentration of GTP $\gamma$ [<sup>35</sup>S] in binding buffer  
170  $\mu$ l membranes from cells expressing neurotensin receptors diluted to 20  $\mu$ g/170  $\mu$ l in binding buffer.

5 **Binding buffer:**

50 mM Hepes/5 mM MgCl<sub>2</sub>, pH 7.4

150 mM NaCl

2  $\mu$ M GDP

1 mM dithiothreitol (DTT)

10 **Protease inhibitors**

100  $\mu$ g/ml bacitracin

100  $\mu$ g/ml benzamidine

5  $\mu$ g/ml aprotinin

5  $\mu$ g/ml leupeptin

15 The protease inhibitors can be obtained from Sigma, St. Louis, MO.

**Wash buffer:**

50 mM Hepes/10 mM MgCl<sub>2</sub>, pH 7.4, ice cold

20 **Procedure:**

1. Set up assay in a 96 well filtering system (Unifilter<sup>®</sup> GF/C<sup>™</sup>, Packard Instrument Company, Meriden, CT).

2. Incubate 60-90 minutes shaking at room temperature.

3. Using a cell Harvester, aspirate samples into processing head. Use a pre-soaked

25 (water is fine) filter. (Do not use polyethylene imine (PEI))

4. Wash four times with cold wash buffer.

5. Dry plate, add 25  $\mu$ l of scintillation fluid to each well.

6. Count samples.

30 **Data Analysis:**

Analyze data as you would for a radioligand binding saturation curve. Subtract the mean of the "minus" wells from the mean of the "plus" wells. Then convert to

Downloaded from www.sciencedirect.com

pmoles/mg protein bound and perform non-linear regression to determine the  $K_d$  for  $\text{GTP}\gamma[^{35}\text{S}]$  binding.

#### $\text{Ca}^{++}$ MOBILIZATION USING THE FLIPR ASSAY

5 Another measure of agonist efficacy and potency is the stimulation of intracellular calcium release. Receptors coupled to the heterotrimeric G protein  $\text{G}_q$  leads to stimulation of phospholipase C. Activation of Phospholipase C leads to the production of diacylglycerol and 1,4,5-inositol-trisphosphate ( $\text{IP}_3$ ).  $\text{IP}_3$  then binds to  $\text{IP}_3$  receptors stimulating release of intracellular  $\text{Ca}^{++}$ . Stimulation of  $\text{Ca}^{++}$  release can be  
10 measured using fluorescent dyes that bind  $\text{Ca}^{++}$  such as Fluo-4AM and Fura-2AM (Molecular Probes, Inc., Eugene, OR). The following is a protocol for measuring intracellular  $\text{Ca}^{++}$  release mediated by neurotensin agonists. The measurement of  $\text{Ca}^{++}$  released upon GPCR activation is known to those skilled in the art and other methods can also be used.

15 All solution components are listed below.

1. Prepare hepes saline and probenecid solutions fresh for each assay. 1 L of assay buffer is sufficient for a small-scale experiment.
- 20 2. Prepare 11 ml dye solution per 96 well cell plate. Aliquot 100  $\mu\text{l}$  per well; incubate at 37°C,  $\text{CO}_2$  incubator, 1 hour.
3. Prepare plate containing compound to be tested, diluting compound to be tested in assay buffer (prepare assay buffer just before use). Allow 100  $\mu\text{l}$  excess compound to be tested volume per well. For example, if 50  $\mu\text{l}$  compound to be  
25 tested will be added to 100  $\mu\text{l}$  cells, then add 150  $\mu\text{l}$  3x concentrated compound per well.
4. Empty media from cell plate and wash with assay buffer on scatron plate washer (if doing large amounts of plates). For smaller amount of plates, can aspirate carefully. Equilibrate cells 10 minutes in assay buffer.
- 30 5. Load cell plate and drug plate into FLIPR; run experiment.

#### **Hepes saline - 1 L**

29 ml 5M NaCl (145 mM final)

- 5 ml 2M Glucose (10 mM final)
- 1.65 ml 3M KCl (5 mM final)
- 1 ml 1M MgSO<sub>4</sub> (1 mM final)
- 10 ml 1M Hepes (10 mM final)
- 5 2 ml 1M CaCl<sub>2</sub> (2 mM final)

**Probenecid Solution - 12.5 ml**

- 925 mg probenecid (2.5 mM final)
- 1.25 ml 5 N NaOH
- 10 11.25 ml Hepes saline

**Dye solution (11 ml)**

1. Combine 11 ml serum-free medium and 110 µl probenecid solution.
2. In a separate tube, add 22 µl dimethylsulfoxide (DMSO) to one vial fluo-4 AM; resuspend by pipetting up and down. Add 22 µl 20% pluronics to this vial. Mix by pipetting--pluronics is very viscous. NOTE: Two fluo-4 vials can be used, and will result in a stronger signal. In this case, double DMSO/pluronics volumes.
3. Add fluo-3/pluronics suspension to serum-free medium; mix well by inversion.

**20 Assay Buffer (1% probenecid solution in hepes saline)**

- 10 ml probenecid solution
- 1 L hepes saline

**25 SATURATION BINDING: [<sup>125</sup>I]NEUROTENSIN (NT) AT NEUROTENSIN RECEPTORS.**

	Well		Final [[ <sup>125</sup> I]NT]
	-	+	
30	A	1,2 3,4	0.02 nM
	B	1,2 3,4	0.05 nM
	C	1,2 3,4	0.75 nM
	D	1,2 3,4	0.10 nM

E	1,2	3,4	0.25 nM
F	1,2	3,4	0.50 nM
G	1,2	3,4	0.75 nM
H	1,2	3,4	1.00 nM

- 5 Total volume in each well is 200  $\mu$ l.

Specific activity of [ $^{125}$ I]NT is 2200 Ci/mmol. Radioactivity calculators that are well known in the art can be used to calculate the specific activities of the stock solutions.

- 10 **To each well add in order:**

20  $\mu$ l buffer to "-" wells

20  $\mu$ l 10  $\mu$ M NT to "+" wells

10  $\mu$ l appropriate concentration of [ $^{125}$ I]NT

170  $\mu$ l membranes diluted to 10  $\mu$ g/170  $\mu$ l

- 15

**Binding buffer:**

50 mM Hepes/10 mM  $MgCl_2$ , pH 7.4

0.2 % BSA (fraction V)

Protease inhibitors

- 20

100  $\mu$ g/ml bacitracin

100  $\mu$ g/ml benzamidine

5  $\mu$ g/ml aprotinin

5  $\mu$ g/ml leupeptin

- 25 **Wash buffer:**

50 mM Hepes/10 mM  $MgCl_2$ , pH 7.4, ice cold.

**Procedure:**

1. Set up assay in a 96 well filtering system (Unifilter<sup>®</sup> GF/C<sup>™</sup>, Packard Instrument Company, Meriden, CT).
2. Incubate 90-120 minutes shaking at room temperature
3. Using a cell Harvester, aspirate samples into processing head. Use a filter pre-soaked in 0.3% PEI.

4. Wash four times with cold wash buffer.
5. Dry plate, add 25  $\mu$ l scintillation fluid to each well.
6. Count samples.

5 Data Analysis

Subtract the mean of the "plus" wells from the mean of the "minus" wells. Then convert to fmoles/mg protein bound and perform non-linear regression to determine the  $K_d$  for [ $^{125}$ I]NT binding.

- 10 Note: This procedure can be modified to use tritiated neurotensin.

COMPETITION BINDING: [ $^{125}$ I]NT AT NEUROTENSIN RECEPTORS

- Up to seven compounds can be tested in 7 point competition curves in a 96 well format. The first six rows for each compound will be used for testing 6 compounds at 6 concentrations in duplicate. An example for a single compound is outlined below. The next compound would be in rows A-F, columns 3 and 4. A seventh compound can be placed in row G1-12 (-9 to -4).

	A1,2	-9
20	B1,2	-8
	C1,2	-7
	D1,2	-6
	E1,2	-5
	F1,2	-4

25

Wells H1,2 are for total counts per minute (cpm) bound.

Wells H3,4 are for 1  $\mu$ M NT to determine non-specific binding.

Filter blanks (just buffer, no membranes) are in H5,6.

- 30 Samples are made in the following stock concentrations:  $10^{-3}$ ,  $10^{-4}$ ,  $10^{-5}$ ,  $10^{-6}$ ,  $10^{-7}$ ,  $10^{-8}$  M. The final concentrations will be one order of magnitude less ( $10^{-4}$  to  $10^{-9}$ ). Stock concentration of compounds are usually 25 mM so a 25:1 dilution is required. Make up 6 tubes labeled -4 to -9. Put 100  $\mu$ l of binding buffer in each tube. Add 4  $\mu$ l

of 25 mM stock to the tube labeled -4. Vortex and take 11  $\mu$ l of the -4 sample and add to the -5 tube. Repeat until all the dilutions are made.

**To each well add in order:**

- 5    20  $\mu$ l buffer to "total" wells (H1, 2).  
     20  $\mu$ l 10  $\mu$ M NT to wells H3, 4.  
     170  $\mu$ l buffer to wells H5, 6.  
     20  $\mu$ l of each concentration of compound to the appropriate wells.  
     10  $\mu$ l of 5 nM [ $^{125}$ I]NT to all wells.
- 10   170  $\mu$ l membranes diluted to 10  $\mu$ g/170  $\mu$ l.

**Procedure:**

1. Set up assay in a 96 well filtering system (Unifilter<sup>®</sup> GF/C<sup>™</sup>, Packard Instrument Company, Meriden, CT).
- 15   2. Incubate 90-120 minutes shaking at room temperature.
3. Using a cell Harvester, aspirate samples into processing head. Use a pre-soaked (0.3% PEI) filter.
4. Wash four times with cold wash buffer.
5. Dry plate, add 25  $\mu$ l of scintillation fluid to each well.
- 20   6. Count samples.

**Binding buffer:**

- 50 mM Hepes/10 mM MgCl<sub>2</sub>, pH 7.4 (Made from 10X stock)  
0.2 % BSA (fraction V)
- 25   Protease inhibitors (Made up as 100X stock).  
     100  $\mu$ g/ml bacitracin  
     100  $\mu$ g/ml benzamidine  
     5  $\mu$ g/ml aprotinin  
     5  $\mu$ g/ml leupeptin

30

**Wash buffer:**

50 mM Hepes/10 mM MgCl<sub>2</sub>, pH 7.4, ice cold (Made from 10X stock).

00000 9271650



# AGONIST-MEDIATED GTP $\gamma$ [<sup>35</sup>S] BINDING ASSAY

Up to seven compounds can be tested in 7 point competition curves in a 96 well format. The first six rows for each compound will be used for testing 6 compounds at 6 concentrations in duplicate. An example for a single compound is outlined below. The next compound would be in rows A-F, columns 3 and 4. A seventh compound can be placed in row G1-12 (-9 to -4).

A1,2 -9  
B1,2 -8  
10 C1,2 -7  
D1,2 -6  
E1,2 -5  
F1,2 -4

15 Wells H1-4 are for basal GTP $\gamma$ [<sup>35</sup>S] bound.  
Wells H5,6 are for cold 10  $\mu$ M GTP $\gamma$ S to determine non-specific binding.  
Filter blanks (just buffer, no membranes) are in H7,8.

20 Samples are made in the following stock concentrations: 10<sup>-3</sup>, 10<sup>-4</sup>, 10<sup>-5</sup>, 10<sup>-6</sup>, 10<sup>-7</sup>, 10<sup>-8</sup> M. The final concentrations will be one order of magnitude less (10<sup>-4</sup> to 10<sup>-9</sup>).  
Stock concentration of compounds are usually 25 mM so a 25:1 dilution is required.  
Make up 6 tubes labeled -4 to -9. Put 100  $\mu$ l of binding buffer in each tube. Add 4  $\mu$ l of 25 mM stock to the tube labeled -4. Vortex and take 11  $\mu$ l of the -4 sample and add to the -5 tube. Repeat until all the dilutions are made. The first sample is always  
25 native neurotensin.

## **To each well add in order:**

20  $\mu$ l binding buffer to wells H1-4.  
20  $\mu$ l of each concentration of compound to the appropriate wells.  
30 20  $\mu$ l of 100  $\mu$ M GTP $\gamma$ S to wells H5,6.  
10  $\mu$ l appropriate concentration of GTP $\gamma$ [<sup>35</sup>S]  
170  $\mu$ l membranes diluted to 20  $\mu$ g/170  $\mu$ l in binding buffer.

**GTP $\gamma$ S binding buffer:**

(Make fresh)

50 mM Hepes/5 mM MgCl<sub>2</sub>, pH 7.4

150 mM NaCl (3M)

5 2  $\mu$ M GDP (10 mM)

1 mM DTT (1 M)

Protease inhibitors

100  $\mu$ g/ml bacitracin

100  $\mu$ g/ml benzamidine

10 5  $\mu$ g/ml aprotinin

5  $\mu$ g/ml leupeptin

**Wash buffer:**

50 mM Hepes/10 mM MgCl<sub>2</sub>, pH 7.4, ice cold

15

**Procedure:**

1. Set up assay in a 96 well filtering system (Unifilter<sup>®</sup> GF/C<sup>™</sup>, Packard Instrument Company, Meriden, CT).

2. Incubate 60 minutes shaking at room temperature.

20 3. Using a cell Harvester, aspirate samples into processing head. Use a pre-soaked (water is fine) filter. Do not use PEI.

4. Wash four times with cold wash buffer.

5. Dry plate, add 25  $\mu$ l scintillation fluid to each well.

1. 6. Count samples.

25

**Data Analysis:**

Analyze data as you would for a dose response curve. Convert to pmoles/mg bound.

30 MEMBRANE PREPARATIONS FROM CELLS

2. Harvest cells. For adherent cells (usually prepare 10 150 mm plates) use PBS/EDTA to remove the cells. Spin down at 1000 X g for 10 minutes at 4° C.

- 5 3. Homogenize with 20 strokes of a tight-fitting glass/glass dounce homogenizer.
4. Spin out nuclei and unlysed cells by centrifuging samples at 1000 X g for 10 minutes at 4° C.
5. Transfer supernatant to new tube.
6. Do two centrifugations at 25,000 X g for 20 minutes in a Sorvall SS34 (or comparable) at 4° C.
- 10 7. Resuspend in an appropriate volume of homogenization buffer that will yield a protein concentration 1-5 mg/ml. Make 250 µl aliquots of each sample plus one 10 µl aliquot for protein determination. For the protein assay dilute 1:5 and measure.

## 15

20

10 mM Na bicarbonate pH 7.4

## 20

25

0.5 mg/ml leupeptin

25

30